

# **Product datasheet for SC209872**

### **OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

Rockville, MD 20850, US
Phone: +1-888-267-4436
https://www.origene.com
techsupport@origene.com
EU: info-de@origene.com
CN: techsupport@origene.cn

## AVPR V2 (AVPR2) (NM\_001146151) Human 3' UTR Clone

#### **Product data:**

**Product Type:** 3' UTR Clones

Product Name: AVPR V2 (AVPR2) (NM\_001146151) Human 3' UTR Clone

**Vector:** pMirTarget (PS100062)

Symbol: AVPR2

Synonyms: ADHR; DI1; DIR; DIR3; NDI; NDI1; V2R

**ACCN:** NM\_001146151

**Insert Size:** 760 bp

Insert Sequence: >SC209872 3'UTR clone of NM\_001146151

The sequence shown below is from the reference sequence of NM\_001146151. The complete

sequence of this clone may contain minor differences, such as SNPs.

Blue=Stop Codon Red=Cloning site

GGCAAGTTGGACGCCCGCAAGATCCGCGAGATTCTCATTAAGGCCAAGAAGGCGGAAAGATCGCCGTG

TAACAATTGGCAGAGCTCAGAATTCAAGCGATCGCC

Restriction Sites: Sgfl-Mlul

**OTI Disclaimer:** Our molecular clone sequence data has been matched to the sequence identifier above as a

point of reference. Note that the complete sequence of this clone is largely the same as the

reference sequence but may contain minor differences, e.g., single nucleotide

polymorphisms (SNPs).





### AVPR V2 (AVPR2) (NM\_001146151) Human 3' UTR Clone - SC209872

**Components:** The cDNA clone is shipped in a 2-D bar-coded Matrix tube as 10 ug dried plasmid DNA. The

package also includes 100 pmols of both the corresponding 5' and 3' vector primers in

separate vials.

**RefSeq:** NM 001146151.3

Summary: This gene encodes the vasopressin receptor, type 2, also known as the V2 receptor, which

belongs to the seven-transmembrane-domain G protein-coupled receptor (GPCR) superfamily, and couples to Gs thus stimulating adenylate cyclase. The subfamily that includes the V2 receptor, the V1a and V1b vasopressin receptors, the oxytocin receptor, and isotocin and mesotocin receptors in non-mammals, is well conserved, though several members signal via other G proteins. All bind similar cyclic nonapeptide hormones. The V2 receptor is expressed in the kidney tubule, predominantly in the distal convoluted tubule and collecting ducts, where its primary property is to respond to the pituitary hormone arginine vasopressin (AVP) by stimulating mechanisms that concentrate the urine and maintain water homeostasis in the organism. When the function of this gene is lost, the disease Nephrogenic Diabetes Insipidus (NDI) results. The V2 receptor is also expressed outside the kidney although its tissue localization is uncertain. When these 'extrarenal receptors' are stimulated by infusion of a V2 selective agonist (dDAVP), a variety of clotting factors are released into the bloodstream. The physiologic importance of this property is not known - its absence does not appear to be detrimental in NDI patients. The gene expression has also been described in

fetal lung tissue and lung cancer associated with alternative splicing. [provided by RefSeq, Jul

2008]

Locus ID: 554

MW: 26.9