

Product datasheet for SC210452

IFIT3 (NM_001031683) Human 3' UTR Clone

Product data:

Product Type: 3' UTR Clones

Symbol: IFIT3

Synonyms: CIG-49; cig41; GARG-49; IFI60; IFIT4; IRG2; ISG60; P60; RIG-G

Mammalian Cell Neomycin

Selection:

Vector: pMirTarget (PS100062)

ACCN: NM_001031683

Insert Size: 870 bp

Insert Sequence: >SC210452 3'UTR clone of NM_001031683

The sequence shown below is from the reference sequence of NM_001031683. The complete sequence

of this clone may contain minor differences, such as SNPs.

Blue=Stop Codon Red=Cloning site

GGCAAGTTGGACGCCCGCAAGATCCGCGAGATTCTCATTAAGGCCAAGAAGGGCGGAAAGATCGCCGTG

TAACAATTGGCAGAGCTCAGAATTCAAGCGATCGCC

GATAGTAATGCTACAATAAATCCAAACATTTCAACTCTGTTA

ACGCGTAAGCGGCCGCGCATCTAGATTCGAAGAAAATGACCGACCAAGCGACGCCCAACCTGCCATCA

CGAGATTTCGATTCCACCGCCGCCTTCTATGAAAGG

Restriction Sites: Safl-Mlul



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com

EU: info-de@origene.com CN: techsupport@origene.cn



IFIT3 (NM_001031683) Human 3' UTR Clone | SC210452

Our molecular clone sequence data has been matched to the sequence identifier above as a

point of reference. Note that the complete sequence of this clone is largely the same as the reference sequence but may contain minor differences, e.g., single nucleotide polymorphisms

(SNPs).

Components: The cDNA clone is shipped in a 2-D bar-coded Matrix tube as 10 ug dried plasmid DNA. The

package also includes 100 pmols of both the corresponding 5' and 3' vector primers in

separate vials.

Note: Plasmids are not sterile. For experiments where strict sterility is required, filtration with 0.22um

filter is required.

RefSeq: <u>NM_001031683.4</u>

Summary: IFN-induced antiviral protein which acts as an inhibitor of cellular as well as viral processes,

cell migration, proliferation, signaling, and viral replication. Enhances MAVS-mediated host antiviral responses by serving as an adapter bridging TBK1 to MAVS which leads to the activation of TBK1 and phosphorylation of IRF3 and phosphorylated IRF3 translocates into nucleus to promote antiviral gene transcription. Exihibits an antiproliferative activity via the up-regulation of cell cycle negative regulators CDKN1A/p21 and CDKN1B/p27. Normally, CDKN1B/p27 turnover is regulated by COPS5, which binds CDKN1B/p27 in the nucleus and

exports it to the cytoplasm for ubiquitin-dependent degradation. IFIT3 sequesters COPS5 in the cytoplasm, thereby increasing nuclear CDKNIB/p27 protein levels. Upregulates CDKNIA/p21 by downregulating MYC, a repressor of CDKNIA/p21. Can negatively regulate the apoptotic effects

of IFIT2.[UniProtKB/Swiss-Prot Function]

Locus ID: 3437

MW: 32.2