

Product datasheet for SC210708

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

MCK10 (DDR1) (NM_013994) Human 3' UTR Clone

Product data:

Product Type: 3' UTR Clones

Product Name: MCK10 (DDR1) (NM 013994) Human 3' UTR Clone

Symbol: MCK10

Synonyms: CAK; CD167; DDR; EDDR1; HGK2; MCK10; NEP; NTRK4; PTK3; PTK3A; RTK6; TRKE

Mammalian Cell

Selection:

Neomycin

Vector: pMirTarget (PS100062)

ACCN: NM_013994

Insert Size: 890 bp

Insert Sequence: >SC210708 3'UTR clone of NM_013994

The sequence shown below is from the reference sequence of NM_013994. The complete

sequence of this clone may contain minor differences, such as SNPs.

Blue=Stop Codon Red=Cloning site

GGCAAGTTGGACGCCCGCAAGATCCGCGAGATTCTCATTAAGGCCAAGAAGGGCGGAAAGATCGCCGTG

 ${\sf TAACAATTGGCAGAGCTCAGAATTCAAGCGATCGCC}$

CGAGATTTCGATTCCACCGCCGCCTTCTATGAAAGG

Restriction Sites: Sgfl-Mlul





MCK10 (DDR1) (NM_013994) Human 3' UTR Clone - SC210708

OTI Disclaimer: Our molecular clone sequence data has been matched to the sequence identifier above as a

point of reference. Note that the complete sequence of this clone is largely the same as the

reference sequence but may contain minor differences, e.g., single nucleotide

polymorphisms (SNPs).

Components: The cDNA clone is shipped in a 2-D bar-coded Matrix tube as 10 ug dried plasmid DNA. The

package also includes 100 pmols of both the corresponding 5' and 3' vector primers in

separate vials.

RefSeq: <u>NM 013994.3</u>

Summary: Receptor tyrosine kinases play a key role in the communication of cells with their

microenvironment. These kinases are involved in the regulation of cell growth, differentiation and metabolism. The protein encoded by this gene belongs to a subfamily of tyrosine kinase receptors with homology to Dictyostelium discoideum protein discoidin I in their extracellular domain, and that are activated by various types of collagen. Expression of this protein is restricted to epithelial cells, particularly in the kidney, lung, gastrointestinal tract, and brain. In addition, it has been shown to be significantly overexpressed in several human tumors. Alternatively spliced transcript variants encoding different isoforms have been described for

this gene. [provided by RefSeq, Feb 2011]

Locus ID: 780 MW: 32