

Product datasheet for SC212084

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Cyclin D3 (CCND3) (NM_001136125) Human 3' UTR Clone

Product data:

Product Type: 3' UTR Clones

Product Name: Cyclin D3 (CCND3) (NM_001136125) Human 3' UTR Clone

Vector: pMirTarget (PS100062)

Symbol: CCND3

ACCN: NM 001136125

Insert Size: 1018 bp

Insert Sequence: >SC212084 3'UTR clone of NM_001136125

The sequence shown below is from the reference sequence of NM_001136125. The complete

sequence of this clone may contain minor differences, such as SNPs.

Blue=Stop Codon Red=Cloning site

GGCAAGTTGGACGCCCGCAAGATCCGCGAGATTCTCATTAAGGCCAAGAAGGGCGGAAAGATCGCCGTG

TAACAATTGGCAGAGCTCAGAATTCAAGCGATCGCC

TGGTGGCCCAGGACCATCATCCTACTGTAATAAAGATGATTGTGAAATAAAA

CGAGATTTCGATTCCACCGCCGCCTTCTATGAAAGG

Restriction Sites: Sgfl-Mlul

OTI Disclaimer: Our molecular clone sequence data has been matched to the sequence identifier above as a

point of reference. Note that the complete sequence of this clone is largely the same as the

reference sequence but may contain minor differences, e.g., single nucleotide

polymorphisms (SNPs).





Cyclin D3 (CCND3) (NM_001136125) Human 3' UTR Clone - SC212084

Components: The cDNA clone is shipped in a 2-D bar-coded Matrix tube as 10 ug dried plasmid DNA. The

package also includes 100 pmols of both the corresponding 5' and 3' vector primers in

separate vials.

RefSeq: <u>NM 001136125.3</u>

Summary: The protein encoded by this gene belongs to the highly conserved cyclin family, whose

members are characterized by a dramatic periodicity in protein abundance through the cell

cycle. Cyclins function as regulators of CDK kinases. Different cyclins exhibit distinct

expression and degradation patterns which contribute to the temporal coordination of each mitotic event. This cyclin forms a complex with and functions as a regulatory subunit of CDK4 or CDK6, whose activity is required for cell cycle G1/S transition. This protein has been shown to interact with and be involved in the phosphorylation of tumor suppressor protein Rb. The

CDK4 activity associated with this cyclin was reported to be necessary for cell cycle progression through G2 phase into mitosis after UV radiation. Several transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Oct 2008]

Locus ID: 896

MW: 36.7