

Product datasheet for **SC213262**

DUSP6 (NM_022652) Human 3' UTR Clone

Product data:

Product Type:	3' UTR Clones
Product Name:	DUSP6 (NM_022652) Human 3' UTR Clone
Symbol:	DUSP6
Synonyms:	HH19; MKP3; PYST1
Mammalian Cell Selection:	Neomycin
Vector:	pMirTarget (PS100062)
ACCN:	NM_022652
Insert Size:	2000 bp

[View online »](#)

Insert Sequence: >SC213262 3'UTR clone of NM_022652
The sequence shown below is from the reference sequence of NM_022652. The complete sequence of this clone may contain minor differences, such as SNPs.
Blue=Stop Codon **Red**=Cloning site

```

GGCAAGTTGGACGCCCGCAAGATCCGCGAGATTCTCATTAAGGCCAAGAAGGGCGGAAAGATCGCCGTG
TAACAATTGGCAGAGCTCAGAATTCAGCGATCGCC
TACCAGGTGGACTCTCTGCAATCTACGGAAAAGACCCACACCCCTCCTTGCTGGAATGTGTCTGGCCC
TTCAGCAGTTTCTCTTGCGAGCATCAGCTGGGCTGCTTTCTTTGTGTGTGGCCCCAGGTGTCAAAATGA
CACCAGCTGTCTGTACTAGACAAGGTTACCAAGTGCAGGAATTGGTTAATACTAACAGAGAGATTTGCTC
CATTCTCTTTGGAATAACAGGACATGCTGTATAGATACAGGCAGTAGGTTTGTCTGTACCCATGTGTA
CAGCCTACCCATGCAGGGACTGGGATTCGAGGACTTCCAGGCGCATAGGGTAGAACCAATGATAGGGT
AGGAGCATGTGTTCTTTAGGGCCTTGTAAGGCTGTTTCCTTTTGCATCTGGAAGTACTATATAATTGT
CTTCAATGAAGACTAATTCAATTTTGCATATAGAGGAGCCAAAGAGAGATTTAGCTCTGTATTTGTGG
TATCAGTTTGAAAAAAAATCTGATACTCCATTTGATTATTGTAATATTTGATCTTGAATCACTTGA
CAGTGTGTTGTTGAATTGTGTTGTTTTCTTTGATGGGCTTAAAGAAATATCCAAAGGGAGAAA
GAGCAGTATGCCACTTCTTAAACAGAACAAAACAAAAAAGAAATTTGTCTCTTTTCTAATCCAAAG
GGTATATTTGCAGCATGCTTGACTTTACCAATTCTGATGACATCTTTACGGACACTATTATCACTAAGA
CCTTGTTATGGCGAAGTCTTTAGTCTTTTTCATGTATTTTCTCATGATTTTTCTCTTTATGTAGTTT
GACTATGCCTTACCTTTGTAAATATTTTGCTTGTTGTGCGAAAGGGGATAATCTGGGAAAGACACC
AAATCATGGGCTCACTTTAAAAAAGAAAGAATAAAAAACCTTCAGCTGTGCTAAACAGTATATTACC
TCTGTATAAAATCTTTCAGGGAGTGTCACTCAATGCAATACTTTGGGTTGGTTTCTTTCTTTAAAA
AAATTTGTATAAACTGGAAGTGTGTGTGTGAGCATGGGTACCCATTTGATAAGAGAAATGCATTG
ATTGTGAAGAAGGGAGAGTTAAATTCTCCATTATGTTTCGTGGTGTAAAGTTTAGAGCTGGAATTTATTA
TAAGAATGTAAACCTTAAATTATTAATAAATACTATTTGGCTATTGAATGTGTGTTTTTAAAACT
CAGTTATACCTATGTGTGAAGTGACACATTTTAGAACTTTTTTATTTCAGGGATTGATTTTGACTGTAG
GCATAGTAGGGCAAGGTTCTTTTAAAGCTTTTCTAGAACGTTTTTCAGCAGCAGTTTATGAAATTCATC
GTTTGCCACGAATAAAGATGACATCCTAGAATGGACCGACTGTACCGTGTATGAACAATACCTCTTA
TGAGTATAGTTCCGGCTTAAGATTTTAAAGCCCTGAAGAAAAATGAACCTATAGAGCATAGCTTGCTG
AGGAATACAATTTGGATATTAACCACAGAGAGGTGTAGCAGAATACCAAGATGACAGCCACATGTTAAC
CTCCTCCCTATCCTTCACTGTCAGGATCTTCTGCTTTATAATGCTGAGATAATGCCAAAGAGGAAGTAG
GGATAATTCCTTTTCCAGTCTGTTGTCATGGGCAATTCTACCTGCAGATATCCTAGTGTGAGAATGCTG
AATCCAGAATGATGTCTCCACTTTTTCCCTAGCATGCGCATGGCAAGGCATATCTAAGCTTCCTCACC
GTGTCAGGATATGGCTTTAGAATTTTACATTGGCACTTAGGAGGCAACTTTGTCATCCCTGTTGTAATG
TGCTTTTTTTGTTAACTGGCTTGGGATGTGGGTGAGAAACCTTATTGAACTTGTGGGGATTATTTTA
GTTTCTTTTCAAGCTAAAGACTTATATCTTGCCATTTTTCTTGGGACCTACCCTCTCCTTACCTCAA
ACGCGTAAGCGGCCGCGGCATCTAGATTGGAAGAAATGACCGACCAAGCGACGCCAACCTGCCATCA
CGAGATTCGATTCCACCGCCGCTTCTATGAAAGG
  
```

Restriction Sites: Sgfl-MluI

OTI Disclaimer: Our molecular clone sequence data has been matched to the sequence identifier above as a point of reference. Note that the complete sequence of this clone is largely the same as the reference sequence but may contain minor differences , e.g., single nucleotide polymorphisms (SNPs).

Components: The cDNA clone is shipped in a 2-D bar-coded Matrix tube as 10 ug dried plasmid DNA. The package also includes 100 pmols of both the corresponding 5' and 3' vector primers in separate vials.

RefSeq: [NM_022652.4](#)

Summary:

The protein encoded by this gene is a member of the dual specificity protein phosphatase subfamily. These phosphatases inactivate their target kinases by dephosphorylating both the phosphoserine/threonine and phosphotyrosine residues. They negatively regulate members of the mitogen-activated protein (MAP) kinase superfamily (MAPK/ERK, SAPK/JNK, p38), which are associated with cellular proliferation and differentiation. Different members of the family of dual specificity phosphatases show distinct substrate specificities for various MAP kinases, different tissue distribution and subcellular localization, and different modes of inducibility of their expression by extracellular stimuli. This gene product inactivates ERK2, is expressed in a variety of tissues with the highest levels in heart and pancreas, and unlike most other members of this family, is localized in the cytoplasm. Mutations in this gene have been associated with congenital hypogonadotropic hypogonadism. Alternatively spliced transcript variants have been found for this gene. [provided by RefSeq, Jan 2014]

Locus ID:

1848

MW:

77.2