

Product datasheet for **SC216945**

LAT2 (SLC7A8) (NM_182728) Human 3' UTR Clone

Product data:

Product Type:	3' UTR Clones
Product Name:	LAT2 (SLC7A8) (NM_182728) Human 3' UTR Clone
Symbol:	LAT2
Synonyms:	LAT2; LPI-PC1
Mammalian Cell Selection:	Neomycin
Vector:	pMirTarget (PS100062)
ACCN:	NM_182728
Insert Size:	1912 bp



[View online »](#)

Insert Sequence: >SC216945 3'UTR clone of NM_182728
The sequence shown below is from the reference sequence of NM_182728. The complete sequence of this clone may contain minor differences, such as SNPs.
Blue=Stop Codon **Red**=Cloning site

```

GGCAAGTTGGACGCCCGCAAGATCCGCGAGATTCTCATTAAGGCCAAGAAGGGCGGAAAGATCGCCGTG
TAACAATTGGCAGAGCTCAGAATTCAGCGATCGCC
AAGGACGTGGCGGGGAGCCCCAGCCCTAGGACCACCATTCCCTGGCTACTCTCTCCTTCTCCCTT
TTTATCTACCTCCCTGCCTTGGTCTGCAACACATGCGAGTACACACACACCCCTCTCTGCTTTT
GTCAGGCAGTGGTAGGACTTTGGTGTGGGTGGTGAAGAAATTGTAACAAAACTGACATTCATACCAA
AGAACCAGCTCTCACCCAGGGTCCATGTCCCAGGCCCACTCCAGTGTGCCACACTCCAGCTGC
TGGAGGAGAGGGGAGATGCCAAGGTGCCCTGCAGGACCTCCCTCCGGGCCACACCCTCAGCTGCCTCT
CAGGAACCGGAGCTCATTACTGCCTTCCCTCCCAGGGAGGCCCTTCAGAGAGGAGAGGCCACAGGAGC
TGCATTGTGGGGGACAGGCTCAAGCAATTCTGTCCCATCAAGGGGTGAGCTGGAGAGACCCAAGACC
CTATCTGTTACACAGGGACCCAAATCCAAGGGGATGCTTCCCTCTGCCCTCTTCTGCCCTCCCCA
TCATACCTGCACCCACCCAGCCAGGGCTCCCTGTCCAGAATTCGGTTCTCTCAGGACGCCAACTCCC
AGAGCTAAGGACCAAGGAGAAGAACAGCCTCTCCACCCCAAGCCAGGCGGTTGAGGAACATATTGAGA
AAGGTTGAGATTGCAGAAACCCAGCCCTGCCCTGCCTCTGCATCCAGCCCCAACATGGTGCCAAAG
CTTCCAGAAGCCAAAAGCTTCTGATTTTAAAGTAGTGGGCATCTCTCTCTAATGACGAAGCTGCTC
AGCAACTCCACCTGCCCGCCGAGGAAGGAGCAGTCCCTGTCTATCCCTGCAGCCACTCCAGCACACC
CGCACACAGCCAGCACCCGCCCCACCGTGCCTTCTCTCTCTGGGCTTGGCTTGGGACCAGGTA
CGAAGGATCCCCAAGCCCTTCAGGCCTGAGATCAGAGCCAGATCAGCCTTAAGTCACCTCCCATCCAAG
AACTTGGCCTAAAAATACTCCCTATTTCTAACCTCAGGACGGATCTGATATTAATGCCTTCCCTGG
GAGGAAGGGTGCTTTCCCTCCCTAGAGGTGCCCATTCATACCCTGGGAGACTGAGGAGAGCATTGG
CTGAAGCCAGTTCTTTCCCATCCATCCCAACTCCAATAATCCCCACTCTCGCAGGTCTCAGTGT
CATGCTGTCTTGGGGCAGGGTGAAAGGGTAGTGGCAGCAGGGCGCCCACTCTGGAGATCTCAAAAAAG
GCCCTCTCTGTGGCTGGCAGCTCTGACCTTCCCTGGGCTTCAAAGGAAGGCTATGGAGTTTGTGT
GGGCCCTGCAACCTTCCAGCCACTCTGCTGCACTAAGGACTTAGGATCCTTTTATCACAATCGGGA
TTCTCTCCCCACCCGAATTCTGTCTGCTTAACTGGAATACACAGGAGCCCTTCTGGCCTGGATGG
TGTCTCCAGCTTCCCCGCCAGCTTGGCCACCCATAGTTGGTGAAGATGCCAAGTTGGTCTGAGTTG
TGACCCCTTCAGAGTAGATGCCCGGCAGGCTGGGGTTGGCCCTGGAGGGTCAGGGGACCATCTTCTTA
TTCCCTCTTTTCTATTCTCCAACCTCCTCCCTCCTCAATTATTTTTTTGTAAGTTGATGCCTTA
CTTTTTGGATAAATATTTTTGAAGCTGGTATTTCTATTTCTTTTGGATTTTTTTAATGTAAGGTTGT
TTGGGGGATGGAGTTAGAACCTTAATGATAATTTCTTTCTTTGGTGTAGGTTTTAGAGATTTGTTTTG
TGGAGAGGTTTTTTCTTTGATGTAATAAAATTTAAATGGAAATGAA
ACGCGTAAGCGGCCGCGGCATCTAGATTCAAGAAAATGACCGACCAAGCGACGCCAACCTGCCATCA
CGAGATTCGATTCCACCGCCGCTTCTATGAAAGG
  
```

Restriction Sites: Sgfl-MluI

OTI Disclaimer: Our molecular clone sequence data has been matched to the sequence identifier above as a point of reference. Note that the complete sequence of this clone is largely the same as the reference sequence but may contain minor differences , e.g., single nucleotide polymorphisms (SNPs).

Components: The cDNA clone is shipped in a 2-D bar-coded Matrix tube as 10 ug dried plasmid DNA. The package also includes 100 pmols of both the corresponding 5' and 3' vector primers in separate vials.

RefSeq: [NM_182728.3](#)

Summary:

Sodium-independent, high-affinity transport of small and large neutral amino acids such as alanine, serine, threonine, cysteine, phenylalanine, tyrosine, leucine, arginine and tryptophan, when associated with SLC3A2/4F2hc. Acts as an amino acid exchanger. Has higher affinity for L-phenylalanine than LAT1 but lower affinity for glutamine and serine. L-alanine is transported at physiological concentrations. Plays a role in basolateral (re)absorption of neutral amino acids. Involved in the uptake of methylmercury (MeHg) when administered as the L-cysteine or D,L-homocysteine complexes, and hence plays a role in metal ion homeostasis and toxicity. Involved in the cellular activity of small molecular weight nitrosothiols, via the stereoselective transport of L-nitrosocysteine (L-CNSO) across the transmembrane. Plays an essential role in the reabsorption of neutral amino acids from the epithelial cells to the bloodstream in the kidney.[UniProtKB/Swiss-Prot Function]

Locus ID:

23428

MW:

70.5