

Product datasheet for **SC300346**

PPAR alpha (PPARA) (NM_001001928) Human Untagged Clone

Product data:

Product Type: Expression Plasmids
Product Name: PPAR alpha (PPARA) (NM_001001928) Human Untagged Clone
Tag: Tag Free
Symbol: PPARA
Synonyms: hPPAR; NR1C1; PPAR; PPAR-alpha; PPARalpha
Mammalian Cell Selection: None
Vector: pCMV6-XL4
E. coli Selection: Ampicillin (100 ug/mL)

Fully Sequenced ORF: >OriGene ORF sequence for NM_001001928 edited
 ATGGTGGACACGAAAGCCCACTCTGCCCCCTCTCCCCACTCGAGGCCGGGATCTAGAG
 AGCCCGTTATCTGAAGAGTTCTGCAAGAAATGGGAAACATCCAAGAGATTTTCGCAATCC
 ATCGGCGAGGATAGTTCTGGAAGCTTTGGCTTTACGGAATACCAGTATTTAGGAAGCTGT
 CCTGGCTCAGATGGCTCGGTCATCACGGACACGCTTTCACCAGCTTCGAGCCCTCCTCG
 GTGACTTATCCTGTGGTCCCCGGCAGCGTGGACGAGTCTCCAGTGGAGCATTGAACATC
 GAATGTAGAATCTGCGGGGACAAGGCCCTCAGGCTATCATTACGGAGTCCACGCGTGTGAA
 GGCTGCAAGGGCTTTCTTCGGCGAACGATTCGACTCAAGCTGGTGTATGACAAGTGCAC
 CGCAGCTGCAAGATCCAGAAAAAGAACAGAAACAAATGCCAGTATTGTCGATTTACAAG
 TGCCTTTCTGTCGGGATGTCACACAACGCGATTGTTTTGGACGAATGCCAAGATCTGAG
 AAAGCAAACTGAAAGCAGAAATTTCTTACCTGTGAACATGACATAGAAGATTCTGAAACT
 GCAGATCTCAAAATCTCTGGCCAAGAGAATCTACGAGGCCACTTGAAGAACTTCAACATG
 AACAAAGTCAAAGCCCGGTTCATCTCTCAGGAAAGGCCAGTAACAATCCACCTTTTGTG
 ATACATGATATGGAGACTGTGTATGGCTGAGAAGACGCTGGTGGCCAAGCTGGTGGCC
 AATGGCATCCAGAACAAGGAGGCCGAGGTCCGCATCTTTCACTGCTGCCAGTGCACGTCA
 GTGGAGACCGTCACGGAGCTCACGGAATTCGCCAAGGCCATCCCAGGCTTCGCAAACTTG
 GACCTGAACGATCAAGTGACATTGCTAAAATACGGAGTTTATGAGGCCATATTCGCCATG
 CTGTCTTCTGTGATGAACAAAGACGGGATGCTGGTAGCGTATGGAATGGGTTTATAACT
 CGTGAATTCCTAAAAAGCCTAAGGAAACCGTTCTGTGATATCATGGAACCCAAGTTTGAT
 TTTGCCATGAAGTTCAATGCACTGGAAGTGGATGACAGTATATCTCCCTTTTTGTGGCT
 GCTATCATTTGCTGTGGAGATCGTCCTGGCCTTCTAAACGTAGGACACATTGAAAAAATG
 CAGGAGGGTATTGTACATGTGCTCAGACTCCACCTGCAGAGCAACCACCCGGACGATATC
 TTTCTCTCCAAAACCTTCTCAAAAAATGGCAGACCTCCGGCAGCTGGTACGGAGCAT
 GCGCAGCTGGTGCAGATCATCAAGAAAGACGGAGTCGGATGCTGCGCTGCACCCGCTACTG
 CAGGAGATCTACAGGGACATGTACTGA

Restriction Sites: Please inquire



[View online »](#)

ACCN:	NM_001001928
Insert Size:	1400 bp
OTI Disclaimer:	<p>Due to the inherent nature of this plasmid, standard methods to replicate additional amounts of DNA in E. coli are highly likely to result in mutations and/or rearrangements. Therefore, OriGene does not guarantee the capability to replicate this plasmid DNA. Additional amounts of DNA can be purchased from OriGene with batch-specific, full-sequence verification at a reduced cost. Please contact our customer care team at custsupport@origene.com or by calling 301.340.3188 option 3 for pricing and delivery.</p> <p>The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. More info</p>
OTI Annotation:	The ORF of this clone has been fully sequenced and found to be a perfect match to NM_001001928.2.
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	<ol style="list-style-type: none"> 1. Centrifuge at 5,000xg for 5min. 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA. 3. Close the tube and incubate for 10 minutes at room temperature. 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom. 5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
RefSeq:	NM_001001928.2 , NP_001001928.1
RefSeq Size:	9965 bp
RefSeq ORF:	1407 bp
Locus ID:	5465
UniProt ID:	Q07869
Cytogenetics:	22q13.31
Protein Families:	Druggable Genome, Nuclear Hormone Receptor, Transcription Factors
Protein Pathways:	Adipocytokine signaling pathway, PPAR signaling pathway

Gene Summary:

Peroxisome proliferators include hypolipidemic drugs, herbicides, leukotriene antagonists, and plasticizers; this term arises because they induce an increase in the size and number of peroxisomes. Peroxisomes are subcellular organelles found in plants and animals that contain enzymes for respiration and for cholesterol and lipid metabolism. The action of peroxisome proliferators is thought to be mediated via specific receptors, called PPARs, which belong to the steroid hormone receptor superfamily. PPARs affect the expression of target genes involved in cell proliferation, cell differentiation and in immune and inflammation responses. Three closely related subtypes (alpha, beta/delta, and gamma) have been identified. This gene encodes the subtype PPAR-alpha, which is a nuclear transcription factor. Multiple alternatively spliced transcript variants have been described for this gene, although the full-length nature of only two has been determined. [provided by RefSeq, Jul 2008]

Transcript Variant: This variant (3) differs in the 5' UTR compared to variant 5. Variants 3 and 5 encode the same protein.