

Product datasheet for SC300414

ntt tec

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

OriGene Technologies, Inc.

Apc11 (ANAPC11) (NM_001002247) Human Untagged Clone

Product data:

Product Type: Expression Plasmids

Product Name: Apc11 (ANAPC11) (NM_001002247) Human Untagged Clone

Tag: Tag Free
Symbol: ANAPC11

Synonyms: APC11; Apc11p; HSPC214

Mammalian Cell Neomycin

Selection:

Vector:pCMV6-Entry (PS100001)E. coli Selection:Kanamycin (25 ug/mL)

Fully Sequenced ORF: >SC300414 representing NM_001002247.

Blue=Insert sequence Red=Cloning site Green=Tag(s)

GATCCGGTACCGAGGAGATCTGCCGCCGCGATCGCC

CAGCAGCACTGCCCCATGTGCCGCCAGGAATGGAAGTTCAAGGAG<mark>TGA</mark>


ACGCGTACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGAT

TACAAGGATGACGACGATAAGGTTTAAACGGCCGGC

Restriction Sites: Sgfl-Mlul



Plasmid Map:



ACCN: NM_001002247

Insert Size: 255 bp

OTI Disclaimer: Our molecular clone sequence data has been matched to the reference identifier above as a

point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative

RNA splicing form or single nucleotide polymorphism (SNP).

OTI Annotation: This TrueClone is provided through our Custom Cloning Process that includes sub-cloning

into OriGene's pCMV6 vector and full sequencing to provide a non-variant match to the expected reference without frameshifts, and is delivered as lyophilized plasmid DNA.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).



Reconstitution Method:

- 1. Centrifuge at 5,000xg for 5min.
- 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
- 3. Close the tube and incubate for 10 minutes at room temperature.
- 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
- 5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

RefSeq: NM 001002247.2

RefSeq Size: 863 bp
RefSeq ORF: 255 bp
Locus ID: 51529
UniProt ID: Q9NYG5
Cytogenetics: 17q25.3

Protein Families: Druggable Genome

Protein Pathways: Cell cycle, Oocyte meiosis, Progesterone-mediated oocyte maturation, Ubiquitin mediated

proteolysis

MW: 9.8 kDa

Gene Summary: Together with the cullin protein ANAPC2, constitutes the catalytic component of the anaphase

promoting complex/cyclosome (APC/C), a cell cycle-regulated E3 ubiquitin ligase that controls progression through mitosis and the G1 phase of the cell cycle. The APC/C complex acts by mediating ubiquitination and subsequent degradation of target proteins: it mainly mediates the formation of 'Lys-11'-linked polyubiquitin chains and, to a lower extent, the formation of 'Lys-48'- and 'Lys-63'-linked polyubiquitin chains. May recruit the E2 ubiquitin-conjugating

enzymes to the complex.[UniProtKB/Swiss-Prot Function]

Transcript Variant: This variant (5) uses an alternate splice site in the 5' UTR and lacks an exon in the 3' coding region, which results in a frameshift, compared to variant 1. The encoded isoform (2) has a distinct C-terminus and is shorter than isoform 1. Variants 2, 3, 4, 5, 6, 7, 8, 9,

10, and 11 encode the isoform 2.