

# **Product datasheet for SC300999**

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## SAP155 (SF3B1) (NM\_001005526) Human Untagged Clone

### **Product data:**

**Product Type:** Expression Plasmids

**Product Name:** SAP155 (SF3B1) (NM\_001005526) Human Untagged Clone

Tag: Tag Free Symbol: SAP155

Synonyms: Hsh155; MDS; PRP10; PRPF10; SAP155; SF3b155

Mammalian Cell

Selection:

Neomycin

Vector:pCMV6-Entry (PS100001)E. coli Selection:Kanamycin (25 ug/mL)

**Restriction Sites:** Sgfl-Mlul

**ACCN:** NM\_001005526

**Insert Size:** 435 bp

**OTI Disclaimer:** Due to the inherent nature of this plasmid, standard methods to replicate additional

amounts of DNA in E. coli are highly likely to result in mutations and/or rearrangements. Therefore, OriGene does not guarantee the capability to replicate this plasmid DNA.

Additional amounts of DNA can be purchased from OriGene with batch-specific, full-sequence

verification at a reduced cost. Please contact our customer care team at

custsupport@origene.com or by calling 301.340.3188 option 3 for pricing and delivery.

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This TrueClone is provided through our Custom Cloning Process that includes sub-cloning

into OriGene's pCMV6 vector and full sequencing to provide a non-variant match to the expected reference without frameshifts, and is delivered as lyophilized plasmid DNA.

**Components:** The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).





#### **Reconstitution Method:**

- 1. Centrifuge at 5,000xg for 5min.
- 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
- 3. Close the tube and incubate for 10 minutes at room temperature.
- 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

RefSeq: <u>NM 001005526.2</u>, <u>NP 001005526.1</u>

RefSeq Size: 665 bp
RefSeq ORF: 435 bp
Locus ID: 23451
Cytogenetics: 2q33.1

**Protein Pathways:** Spliceosome

**Gene Summary:** This gene encodes subunit 1 of the splicing factor 3b protein complex. Splicing factor 3b,

together with splicing factor 3a and a 12S RNA unit, forms the U2 small nuclear

ribonucleoproteins complex (U2 snRNP). The splicing factor 3b/3a complex binds pre-mRNA upstream of the intron's branch site in a sequence independent manner and may anchor the U2 snRNP to the pre-mRNA. Splicing factor 3b is also a component of the minor U12-type spliceosome. The carboxy-terminal two-thirds of subunit 1 have 22 non-identical, tandem HEAT repeats that form rod-like, helical structures. Alternative splicing results in multiple

transcript variants encoding different isoforms. [provided by RefSeq, Jul 2008]

Transcript Variant: This variant (2) uses an alternate splice site in the 3' coding region,

compared to variant 1, that results in a frameshift. It encodes a shorter isoform (2) which has

a distinct C-terminus compared to isoform 1.