

Product datasheet for SC301006

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

EDA (NM_001005612) Human Untagged Clone

Product data:

Product Type: Expression Plasmids

Product Name: EDA (NM_001005612) Human Untagged Clone

Tag: Tag Free Symbol: EDA

Synonyms: ECTD1; ED1-A1; ED1-A2; EDA-A1; EDA-A2; EDA1; EDA2; HED1; HED1; ODT1; STHAGX1;

TNLG7C; XHED; XLHED

Vector: pCMV6 series

Fully Sequenced ORF: >NCBI ORF sequence for NM_001005612, the custom clone sequence may differ by one or

more nucleotides

AGCCTGTGA

Restriction Sites: Please inquire **ACCN:** NM_001005612

OTI Disclaimer: Our molecular clone sequence data has been matched to the reference identifier above as a

point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative

RNA splicing form or single nucleotide polymorphism (SNP).

OTI Annotation: This TrueClone is provided through our Custom Cloning Process that includes sub-cloning

into OriGene's pCMV6 vector and full sequencing to provide a non-variant match to the expected reference without frameshifts, and is delivered as lyophilized plasmid DNA.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).





Reconstitution Method:

- 1. Centrifuge at 5,000xg for 5min.
- 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
- 3. Close the tube and incubate for 10 minutes at room temperature.
- 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

RefSeg: NM 001005612.1, NP 001005612.1

RefSeq Size: 1140 bp
RefSeq ORF: 429 bp
Locus ID: 1896
UniProt ID: Q92838
Cytogenetics: Xq13.1

Protein Families: Druggable Genome, Secreted Protein, Transmembrane

Protein Pathways: Cytokine-cytokine receptor interaction

Gene Summary: The protein encoded by this gene is a type II membrane protein that can be cleaved by furin

to produce a secreted form. The encoded protein, which belongs to the tumor necrosis factor

family, acts as a homotrimer and may be involved in cell-cell signaling during the

development of ectodermal organs. Defects in this gene are a cause of ectodermal dysplasia, anhidrotic, which is also known as X-linked hypohidrotic ectodermal dysplasia. Several

transcript variants encoding many different isoforms have been found for this gene.

[provided by RefSeq, Jul 2008]

Transcript Variant: This variant (5) uses two alternate in-frame splice sites compared to variant 1. The resulting isoform (5) has the same N- and C-termini but is shorter than isoform 1. Sequence Note: This RefSeq record was created from transcript and genomic sequence data to make the sequence consistent with the reference genome assembly. The genomic coordinates used for the transcript record were based on transcript alignments.