

Product datasheet for SC301304

Septin 2 (SEPT2) (NM_001008491) Human Untagged Clone

Product data:


Product Type:	Expression Plasmids
Product Name:	Septin 2 (SEPT2) (NM_001008491) Human Untagged Clone
Tag:	Tag Free
Symbol:	SEPTIN2
Synonyms:	DIFF6; hNedd5; NEDD-5; NEDD5; Pnutl3; SEPT2
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)
Fully Sequenced ORF:	>SC301304 representing NM_001008491. Blue=Insert sequence Red=Cloning site Green=Tag(s)

```
GCTCGTTTGTAGTGAACCGTCAGAATTTTGTAAACGACTCACTATAGGGCGCCGGGAATTCGTCGACTG
GATCCGGTACCGAGGAGATCTGCCGCCGCGATCGCC
ATGTCTAAGCAACAGCCAACCTCAGTTTATAAATCCAGAAACACCTGGCTATGTTGGATTTGCAAACCTC
CCCAATCAAGTTCACCGAAAATCAGTGAAGAAAGGTTTGTAGTTACACTGATGGTGGTCCGGTGAATCA
GGTCTAGGAAAATCGACTCTCATAAACAGCCTATTCCTAACTGATCTGTACCCAGAAAGAGTCATACCT
GGAGCAGCAGAAAAAATGAAAGAACTGTCCAGATTGAGGCTTCAACTGTTGAAATTGAAGAGCGAGGG
GTCAAGCTACGCCTGACAGTGGTAGATACCCTGGCTATGGTGACGCTATCAACTGCAGAGATTGTTTT
AAGACAATTATCTCTATATTGATGAGCAATTTGAGAGGTACCTGCATGACGAGAGCGGCTTGAACAGG
CGGCACATCATTGATAATAGGGTGCATTGTTGCTTTTACTTTATTTACCTTTTGGACATGGACTTAAG
CCCTTAGATGTGGCGTTTATGAAGGCAATACACAACAAGGTGAATATTGTGCCTGTCATTGCAAAAAGCT
GACACTCTCACCTGAAGGAACGGGAGCGGCTGAAGAAAAGGATTCTGGATGAAATTGAAGAACATAAC
ATCAAAATCTATCACTTACCTGATGCAGAAATCAGATGAAGATGAAGATTTTAAAGAGCAGACTAGACTT
CTCAAGGCTAGCATCCCATTCTCTGTGGTTGGATCCAATCAGTTGATTGAAGCCAAAGGAAAAGAAGGTC
AGAGGCCGCTCTACCCCTGGGGTGTGTGGAAGTGGAGAAGCCAGAGCAATGACTTTCTGAAGCTG
AGAACCATGCTCATCACCACATGCAGGATCTCCAGGAGGTGACCCAGGACCTTCATTATGAAAACCTC
CGTTCTGAGAGACTCAAGAGAGGCGGCAGGAAAGTGGAGAATGAGGACATGAATAAAGACCAGATCTTG
CTGAAAAAAGAAGCTGAGCTCCGCCGATGCAAGAGATGATTGCAAGGATGCAGGCGCAGATGCAGATG
CAGATGCAGGGCGGGGATGGCGATGGCGGGCTCTCGGGCACCACGTGTA
ACGCGTACGCGGCCGCTCGAGCAGAACTCATCTCAGAAGAGGATCTGGCAGCAATGATATCCTGGAT
TACAAGGATGACGACGATAAGGTTTAAACGGCCGGC
```

Restriction Sites: SgfI-MluI



[View online »](#)

Plasmid Map:


ACCN: NM_001008491

Insert Size: 1086 bp

OTI Disclaimer: Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).

OTI Annotation: This TrueClone is provided through our Custom Cloning Process that includes sub-cloning into OriGene's pCMV6 vector and full sequencing to provide a non-variant match to the expected reference without frameshifts, and is delivered as lyophilized plasmid DNA.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method:

1. Centrifuge at 5,000xg for 5min.
2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
3. Close the tube and incubate for 10 minutes at room temperature.
4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

RefSeq: [NM_001008491.2](#)

RefSeq Size: 3726 bp

RefSeq ORF: 1086 bp

Locus ID: 4735

UniProt ID: [Q15019](#)

Cytogenetics: 2q37.3

MW: 41.5 kDa

Gene Summary: Filament-forming cytoskeletal GTPase. Forms a filamentous structure with SEPTIN12, SEPTIN6, SEPTIN2 and probably SEPTIN4 at the sperm annulus which is required for the structural integrity and motility of the sperm tail during postmeiotic differentiation (PubMed:25588830). Required for normal organization of the actin cytoskeleton. Plays a role in the biogenesis of polarized columnar-shaped epithelium by maintaining polyglutamylated microtubules, thus facilitating efficient vesicle transport, and by impeding MAP4 binding to tubulin. Required for the progression through mitosis. Forms a scaffold at the midplane of the mitotic spindle required to maintain CENPE localization at kinetochores and consequently chromosome congression. During anaphase, may be required for chromosome segregation and spindle elongation. Plays a role in ciliogenesis and collective cell movements. In cilia, required for the integrity of the diffusion barrier at the base of the primary cilium that prevents diffusion of transmembrane proteins between the cilia and plasma membranes: probably acts by regulating the assembly of the tectonic-like complex (also named B9 complex) by localizing TMEM231 protein. May play a role in the internalization of 2 intracellular microbial pathogens, *Listeria monocytogenes* and *Shigella flexneri*. [UniProtKB/Swiss-Prot Function]
Transcript Variant: This variant (1) represents the longest transcript and encodes isoform a. Variants 1-4, 8-13, and 16-18 encode the same isoform (a).