

## **Product datasheet for SC301684**

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## IL32 (NM\_001012635) Human Untagged Clone

## **Product data:**

**Product Type:** Expression Plasmids

Product Name: IL32 (NM\_001012635) Human Untagged Clone

Tag: Tag Free Symbol: IL32

Synonyms: IL-32alpha; IL-32beta; IL-32delta; IL-32gamma; NK4; TAIF; TAIFa; TAIFb; TAIFc; TAIFd

**Vector:** pCMV6 series

Fully Sequenced ORF: >NCBI ORF sequence for NM\_001012635, the custom clone sequence may differ by one or

more nucleotides

TGCTCTGAACCCCAATCCTCAAAATGA

**Restriction Sites:** Please inquire ACCN: NM\_001012635

**OTI Disclaimer:** Our molecular clone sequence data has been matched to the reference identifier above as a

point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative

RNA splicing form or single nucleotide polymorphism (SNP).

OTI Annotation: This TrueClone is provided through our Custom Cloning Process that includes sub-cloning

into OriGene's pCMV6 vector and full sequencing to provide a non-variant match to the expected reference without frameshifts, and is delivered as lyophilized plasmid DNA.

**Components:** The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).





**Reconstitution Method:** 

- 1. Centrifuge at 5,000xg for 5min.
- 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
- 3. Close the tube and incubate for 10 minutes at room temperature.
- 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
- 5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

**RefSeq:** <u>NM 001012635.1</u>, <u>NP 001012653.1</u>

 RefSeq Size:
 892 bp

 RefSeq ORF:
 507 bp

 Locus ID:
 9235

 UniProt ID:
 P24001

 Cytogenetics:
 16p13.3

**Protein Families:** Secreted Protein

**Gene Summary:** This gene encodes a member of the cytokine family. The protein contains a tyrosine sulfation

site, 3 potential N-myristoylation sites, multiple putative phosphorylation sites, and an RGD cell-attachment sequence. Expression of this protein is increased after the activation of T-cells by mitogens or the activation of NK cells by IL-2. This protein induces the production of TNFalpha from macrophage cells. Alternate transcriptional splice variants, encoding different

isoforms, have been characterized. [provided by RefSeq, Jul 2008]

Transcript Variant: This variant (6) differs in the 5' UTR and lacks an alternate in-frame exon within the coding region, compared to variant 1, resulting in a shorter protein (isoform C). Sequence Note: The RefSeq transcript and protein were derived from genomic sequence to

make the sequence consistent with the reference genome assembly. The genomic

coordinates used for the transcript record were based on alignments.