

Product datasheet for **SC302099**

P4HA2 (NM_001017974) Human Untagged Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	P4HA2 (NM_001017974) Human Untagged Clone
Tag:	Tag Free
Symbol:	P4HA2
Synonyms:	MYP25
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)



[View online »](#)

Fully Sequenced ORF: >SC302099 representing NM_001017974.
 Blue=Insert sequence Red=Cloning site Green=Tag(s)

```

GCTCGTTTAGTGAACCGTCAGAATTTTGTAAACGACTCACTATAGGGCGGCCGGGAATTCGTGACTG
GATCCGGTACCGAGGAGATCTGCCGCCCGGATCGCC
ATGAAACTCTGGGTGTCTGCATTGCTGATGGCCTGGTTTGGTGTCTGAGCTGTGTGCAGGCCGAATTC
TTCACCTCTATTGGGCACATGACTGACCTGATTTATGCAGAGAAAGAGCTGGTGCAGTCTCTGAAAGAG
TACATCCTTGTGGAGGAAGCCAAGCTTTCCAAGATTAAGAGCTGGGCCAACAAAATGGAAGCCTTGACT
AGCAAGTCAGCTGCTGATGCTGAGGGCTACCTGGCTCACCTGTGAATGCCTACAACTGGTGAAGCGG
CTAAACACAGACTGGCCTGCGCTGGAGGACCTTGTCTGCAGGACTCAGCTGCAGTTTTATCGCCAAC
CTCTCTGTGCAGCGGCAGTTCTTCCCCTGATGAGGACGAGATAGGAGCTGCCAAAGCCCTGATGAGA
CTTCAGGACACATACAGGCTGGACCCAGGCACAATTTCCAGAGGGGAACCTCCAGGAACCAAGTACCAG
GCAATGCTGAGTGGATGACTGCTTTGGGATGGGCCGCTCGGCCTACAATGAAGGGGACTATTATCAT
ACGGTGTGTGGATGGAGCAGGTGCTAAAGCAGCTTGATGCCGGGAGGAGGCCACCACAACCAAGTCA
CAGGTGCTGGACTACCTCAGCTATGCTGTCTTCCAGTTGGGTGATCTGCACCGTGCCCTGGAGCTCACC
CGCCGCTGCTCTCCCTTGACCCAAGCCACGAACGAGCTGGAGGGAATCTGCGGTACTTTGAGCAGTTA
TTGGAGGAAGAGAGAGAAAAACGTTAACAAATCAGACAGAAGCTGAGCTAGCAACCCAGAAAGGCATC
TATGAGAGGCCTGTGGACTACCTGCCTGAGAGGGATGTTTACGAGAGCCTCTGTCGTGGGGAGGGTGTG
AAACTGACACCCCGTAGACAGAAGAGGCTTTTCTGTAGGTACCACCATGGCAACAGGGCCCCACAGCTG
CTCATTGCCCCCTTCAAAGAGGAGGACGAGTGGGACAGCCCGCACATCGTCAGGTACTACGATGTCATG
TCTGATGAGGAAATCGAGAGGATCAAGGAGATCGCAAAACCTAACTTGCACGAGCCACCGTTCTGAT
CCCAAGACAGGAGTCTCACTGTCGCCAGCTACCGGTTTCCAAAAGCTCCTGGCTAGAGGAAGATGAT
GACCCTGTTGTGGCCCGAGTAAATCGTCGGATGCAGCATATCACAGGGTTAACAGTAAAGATGCAGAA
TTGTTACAGTTGCAAAATTATGGAGTGGGAGGACAGTATGAACCGCACTTCGACTTCTCTAGGCGACCT
TTTGACAGCGGCCTCAAAACAGAGGGGAATAGGTTAGCGACGTTTCTTAACTACATGAGTGTAGAA
GCTGGTGGTGCCACCGTCTTCCCTGATCTGGGGCTGCAATTTGGCCTAAGAAGGGTACAGCTGTGTTT
TGGTACAACCTCTTGGGAGCGGGGAAGGTGACTACCGAACAAGACATGCTGCCTGCCCTGTGCTTGTG
GGCTGCAAGTGGTCTCCAATAAGTGGTTCCATGAACGAGGACAGGAGTCTTGAGACCTTGTGGATCA
ACAGAAGTTGACTGA
ACGGGTACGCGGCCGCTCGAGCAGAACTCATCTCAGAAGAGGATCTGGCAGCAATGATATCTGGAT
TACAAGGATGACGACGATAAGGTTTAAACGGCCGCGC
  
```

Restriction Sites: SgfI-MluI

ACCN: NM_001017974

Insert Size: 1602 bp

OTI Disclaimer: Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).

OTI Annotation: This TrueClone is provided through our Custom Cloning Process that includes sub-cloning into OriGene's pCMV6 vector and full sequencing to provide a non-variant match to the expected reference without frameshifts, and is delivered as lyophilized plasmid DNA.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method:	<ol style="list-style-type: none">1. Centrifuge at 5,000xg for 5min.2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.3. Close the tube and incubate for 10 minutes at room temperature.4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
RefSeq:	NM_001017974.1
RefSeq Size:	2110 bp
RefSeq ORF:	1602 bp
Locus ID:	8974
UniProt ID:	O15460
Cytogenetics:	5q31.1
Protein Families:	Druggable Genome
Protein Pathways:	Arginine and proline metabolism, Metabolic pathways
MW:	60.6 kDa
Gene Summary:	<p>This gene encodes a component of prolyl 4-hydroxylase, a key enzyme in collagen synthesis composed of two identical alpha subunits and two beta subunits. The encoded protein is one of several different types of alpha subunits and provides the major part of the catalytic site of the active enzyme. In collagen and related proteins, prolyl 4-hydroxylase catalyzes the formation of 4-hydroxyproline that is essential to the proper three-dimensional folding of newly synthesized procollagen chains. Alternatively spliced transcript variants encoding different isoforms have been described. [provided by RefSeq, Jul 2008]</p> <p>Transcript Variant: This variant (3) differs in the 5' UTR, compared to variant 1, and contains an alternate in-frame exon. Variants 2-4 and 7-9 all encode isoform 2, which is shorter than isoform 1.</p>