

## Product datasheet for SC302126

### EMA (MUC1) (NM\_001018017) Human Untagged Clone

#### Product data:

Product Type:	Expression Plasmids
Product Name:	EMA (MUC1) (NM_001018017) Human Untagged Clone
Tag:	Tag Free
Symbol:	EMA
Synonyms:	ADMCKD; ADMCKD1; ADTKD2; CA 15-3; CD227; EMA; H23AG; KL-6; MAM6; MCD; MCKD; MCKD1; MUC-1; MUC-1/SEC; MUC-1/X; MUC1/ZD; PEM; PEMT; PUM
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)
Fully Sequenced ORF:	>SC302126 representing NM_001018017. Blue=Insert sequence Red=Cloning site Green=Tag(s)

GCTCGTTTAGTGAACCGTCAGAATTTTGTAAACGACTCACTATAGGGCGGCCGGAATTCGTCGACTG  
 GATCCGGTACCGAGGAGATCTGCCGCCGCGATCGCC  
 ATGACACCGGGCACCCAGTCTCCTTTCTTCTGCTGCTCCTCACAGTGCTTACAGTTGTTACGGGT  
 TCTGGTCATGCAAGCTCTACCCAGGTGGAGAAAAGGAGACTTCGGCTACCCAGAGAAGTTCAGTGCCC  
 AGCTCTACTGAGAAGAATGCTTTTAATTCCTCTCTGGAAGATCCAGCACCGACTACTACCAAGAGCTG  
 CAGAGAGACATTTCTGAAATGTTTTGAGATTATAAACAAGGGGGTTTTCTGGGCCTCTCCAATATT  
 AAGTTCAGGCCAGGATCTGTGGTGTACAATTGACTCTGGCCTTCGAGAAGGTACCATCAATGTCCAC  
 GACGTGGAGACACAGTTCAATCAGTATAAACGGAAGCAGCCTCTCGATATAACCTGACGATCTCAGAC  
 GTCAGCGTGAGTGATGTGCCATTTCTTTCTGCCCAGTCTGGGGCTGGGTGCCAGGCTGGGCATC  
 GCGCTGCTGGTGTGTGTCTGTGTTCTGGTTCGCTGGCCATTGTCTATCTCATTGCCTTGCTGTCTGT  
 CAGTGCCGCCGAAAGAACTACGGGCAGCTGGACATCTTTCCAGCCCGGGATACCTACCATCCTATGAGC  
 GAGTACCCACCTACCACACCATGGGCGCTATGTGCCCCCTAGCAGTACCGATCGTAGCCCTATGAG  
 AAGGTTTCTGCAGGAATGGTGGCAGCAGCCTCTTTACACAAACCCAGCAGTGCCAGCCACTTCTGCC  
 AACTTGTAAG  
 ACGCGTACGCGGCCGCTCGAGCAGAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGAT  
 TACAAGGATGACGACGATAAGGTTTAAACGGCCGCGC

Restriction Sites:	SgfI-MluI
ACCN:	NM_001018017
Insert Size:	768 bp


[View online »](#)

<b>OTI Disclaimer:</b>	Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).
<b>OTI Annotation:</b>	This TrueClone is provided through our Custom Cloning Process that includes sub-cloning into OriGene's pCMV6 vector and full sequencing to provide a non-variant match to the expected reference without frameshifts, and is delivered as lyophilized plasmid DNA.
<b>Components:</b>	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
<b>Reconstitution Method:</b>	<ol style="list-style-type: none"> <li>1. Centrifuge at 5,000xg for 5min.</li> <li>2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.</li> <li>3. Close the tube and incubate for 10 minutes at room temperature.</li> <li>4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.</li> <li>5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.</li> </ol>
<b>RefSeq:</b>	<u>NM_001018017.2</u>
<b>RefSeq Size:</b>	1166 bp
<b>RefSeq ORF:</b>	768 bp
<b>Locus ID:</b>	4582
<b>UniProt ID:</b>	<u>P15941</u>
<b>Cytogenetics:</b>	1q22
<b>Protein Families:</b>	Druggable Genome, Secreted Protein, Transmembrane
<b>MW:</b>	27.6 kDa
<b>Gene Summary:</b>	<p>This gene encodes a membrane-bound protein that is a member of the mucin family. Mucins are O-glycosylated proteins that play an essential role in forming protective mucous barriers on epithelial surfaces. These proteins also play a role in intracellular signaling. This protein is expressed on the apical surface of epithelial cells that line the mucosal surfaces of many different tissues including lung, breast stomach and pancreas. This protein is proteolytically cleaved into alpha and beta subunits that form a heterodimeric complex. The N-terminal alpha subunit functions in cell-adhesion and the C-terminal beta subunit is involved in cell signaling. Overexpression, aberrant intracellular localization, and changes in glycosylation of this protein have been associated with carcinomas. This gene is known to contain a highly polymorphic variable number tandem repeats (VNTR) domain. Alternate splicing results in multiple transcript variants.[provided by RefSeq, Feb 2011]</p> <p>Transcript Variant: This variant (3) uses an alternate in-frame splice site, compared to variant 1. The resulting isoform (3) is shorter than isoform 1.</p>