

## Product datasheet for **SC302859**

### **FO XK1 (NM\_001037165) Human Untagged Clone**

#### **Product data:**

Product Type:	Expression Plasmids
Product Name:	FO XK1 (NM_001037165) Human Untagged Clone
Tag:	Tag Free
Symbol:	FO XK1
Synonyms:	FO XK1L
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)




[View online »](#)

Fully Sequenced ORF: >SC302859 representing NM\_001037165.  
Blue=Insert sequence Red=Cloning site Green=Tag(s)

```
CTATAGGGCGCCGGGAATTCGTGCGACTGGATCCGGTACCGAGGAGATCTGCCGCCGCGATCGCCGGCG
CGCCAGATCTCAAGCTTAACTAGCTAGCGGACCGAC
ATGGCCGAAGTCGGCGAGGACAGCGGCCGCCGCCCTGCTCGCGCTGCGCTCGGCGCCCTGCAGCCCA
GTGTGTGCGCCGAGCCGCCGCCGCCCTTCCCCGCGGCCGACCCCCGCGCCCCGCGCAGCCC
CAGCCTCCGCCCCGGCCGCCGCCGCCGCCGCCGCCGCCGCCGCCGCCGCCGCCGCCGCCGCCGCCGCC
TCTCCGGGGGCTCCTCCGGGGTATCCGGGGACTCCGCGGTGCGGGGCGGGCGCCGCCCTGGTGGCC
GCGGGCGCCGCTCGGTACGGCAGAGCCCGGGCCGGCGCTGGCGCGGTGGAGGGCCGCGAGTTCGAG
TTCCTCATGCGCCAGCCAGCGTCACCATCGGCCGCAACTCGTCGAGGGCTCGGTGGACTTGAGCATG
GGCCTGTCCAGCTTCATCTCGCGGCCACCTGCAGCTCAGTTCAGGAGCCGCACTTCTACCTGCGC
TGCTCGCAAGAACGGCGTCTTCGTGGACGGGGCTTCCAGAGACGCGGGCGCCGCCCTGCAGCTG
CCCAAGCAGTGTACCTCCGGTTTCCAGCACGGCCATCAAGATCCAGTTCAGTCGCTCTATCACAAA
GAAGAGGCCCCAGCTCCCCGCTGCGGCCACTGTACCCCAAGATCTCCCCTGAAGATCCACATCCCG
GAGCCGACCTCCGGAGCATGGTCAGCCCCGTCCTCCCGACGGGCACCATCAGTGTCCCAACTCC
TGCCAGCCAGTCCACGGGTGCGGGTCTCCAGTTACCGCTTTGTGCAGAACGTGACCTCGGACCTG
CAGCTGGCAGCAGAGTTTGCAGCAAAGGCCGCTCGGAGCAGCAGGCAGACAGTCTGGAGGAGACAGC
CCCAAGGATGAGTCAAAGCCGCCGTTCTCTACGCGCAGCTGATCGTGCAGGCCATCTCCTCCGCCAG
GACCGGCAGCTGACCCTGAGCGGGATCTACGCCACATCACCAAGCATTACCCCTACTACCGGACGGCC
GACAAAGGCTGGCAGAATTCTATCCGGCACAACCTCTCTTTGAACCGTACTTTATCAAAGTCCCAGT
TCCCAGGAGGAGCCTGGGAAGGGTCTTTTGGGAATAGACCTGCCTCTGAAGCAAGCTCGTGGAA
CAGGCATTCCGGAAACGGAGGCAGAGGGGTGCTCCTGCTTCCGCACCCCTTCCGGCCCTGTCTCTCA
AGGAGCGCTCCAGCTTCGCCACACACCCCGGGTGTATGCCCTCGCTCCGGCGGCTGCAGACCCCA
GAGTGCCTGTCTCGGAGGGTCTCCCCATTCCACACGACCTGAGTTTGGGTCCAAGTTAGCTTCTGTC
CCAGAGTACCGGTATTCCAAAGCGCACCCGGTCCCCCGTCAGCGCCAGCCAGTATGCGCGTG
CCTCCCCGACCGTCCAGCCTCGTGGCAAGCCGTGGCTACATGCCCGCTCCATCGTAACTCACAG
CAGCCCGCGGGCCAGCCATCCACGTGTCAGCAGGCCCCACCCTCACCATGGTCAGGGTGGTCACC
ACATCTGCCAACTCGGCCAACGGATACATCCTCACCAGCCAGGGCGGGCGGGGGCTCCCATGATGCG
GCGGGCGCAGCCGTGCTGGACCTGGGCAGCGAGGCCAGAGCCTGGAGGAGAAACCACCATTGCGTTT
GCCACAATCCCCGCGGTGGTGGAGTATCCAGACGGTGGCCAGCCAGATGGCCCCGGGGTCCCCGGA
CACACGGTCAACATCCTGACGCCGCCACACCCGTGACCTCGGGCAGCACACCTTCCAGTCCGGGCC
GTGACCCAGAACGAAAGCATGCGGTTCCACGAACAGTTTAGCCGGCAACGCTTACGCCCTCACCAGC
CCTTTGACGCTCCTTGGACCCAAGCGAGTTCATCCGCGCCGGTGGTGGTCAACCGGGTGTGCGAGGTG
GGGCCAAGGAGCCAGCAGCAGCCGTGCGGCCACGGCCACCACCACCCAGCCACTGCCACCACCGCC
TCTGCCTCCGCTCTTCCACTGGAGAGCCCGAGGTCAAAGGTCCCGGGTGGAGGAGCCAGTGGGGCT
GTAACCACACCGGCTGGAGTGTGTCAGCTGCCGGCCCCAGGGCCAGGCACCGGGGAGTGA
ACGGTACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGAT
TACAAGGATGACGACGATAAGGTTTAAACGGCCGGC
```

Restriction Sites: RsrII-MluI

## Plasmid Map:



ACCN: NM\_001037165

Insert Size: 2202 bp

**OTI Disclaimer:** Due to the inherent nature of this plasmid, standard methods to replicate additional amounts of DNA in *E. coli* are highly likely to result in mutations and/or rearrangements. Therefore, OriGene does not guarantee the capability to replicate this plasmid DNA. Additional amounts of DNA can be purchased from OriGene with batch-specific, full-sequence verification at a reduced cost. Please contact our customer care team at [custsupport@origene.com](mailto:custsupport@origene.com) or by calling 301.340.3188 option 3 for pricing and delivery.

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. [More info](#)

**OTI Annotation:** This TrueClone is provided through our Custom Cloning Process that includes sub-cloning into OriGene's pCMV6 vector and full sequencing to provide a non-variant match to the expected reference without frameshifts, and is delivered as lyophilized plasmid DNA.

**Components:** The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

**Reconstitution Method:**

1. Centrifuge at 5,000xg for 5min.
2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
3. Close the tube and incubate for 10 minutes at room temperature.
4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

RefSeq: [NM\\_001037165.1](#)

RefSeq Size: 11191 bp

RefSeq ORF: 2202 bp

Locus ID: 221937

UniProt ID: [P85037](#)

Cytogenetics: 7p22.1

MW: 75.5 kDa

**Gene Summary:** Transcriptional regulator involved in different processes such as glucose metabolism, aerobic glycolysis, muscle cell differentiation and autophagy (By similarity). Recognizes and binds the forkhead DNA sequence motif (5'-GTAAACA-3') and can both act as a transcription activator or repressor, depending on the context (PubMed:17670796). Together with FOXK2, acts as a key regulator of metabolic reprogramming towards aerobic glycolysis, a process in which glucose is converted to lactate in the presence of oxygen (By similarity). Acts by promoting expression of enzymes for glycolysis (such as hexokinase-2 (HK2), phosphofructokinase, pyruvate kinase (PKLR) and lactate dehydrogenase), while suppressing further oxidation of pyruvate in the mitochondria by up-regulating pyruvate dehydrogenase kinases PDK1 and PDK4 (By similarity). Probably plays a role in gluconeogenesis during overnight fasting, when lactate from white adipose tissue and muscle is the main substrate (By similarity). Involved in mTORC1-mediated metabolic reprogramming: in response to mTORC1 signaling, translocates into the nucleus and regulates the expression of genes associated with glycolysis and downstream anabolic pathways, such as HIF1A, thereby regulating glucose metabolism (By similarity). Together with FOXK2, acts as a negative regulator of autophagy in skeletal muscle: in response to starvation, enters the nucleus, binds the promoters of autophagy genes and represses their expression, preventing proteolysis of skeletal muscle proteins (By similarity). Acts as a transcriptional regulator of the myogenic progenitor cell population in skeletal muscle (By similarity). Binds to the upstream enhancer region (CCAC box) of myoglobin (MB) gene, regulating the myogenic progenitor cell population (By similarity). Promotes muscle progenitor cell proliferation by repressing the transcriptional activity of FOXO4, thereby inhibiting myogenic differentiation (By similarity). Involved in remodeling processes of adult muscles that occur in response to physiological stimuli (By similarity). Required to correct temporal orchestration of molecular and cellular events necessary for muscle repair (By similarity). Represses myogenic differentiation by inhibiting MEFC activity (By similarity). Positively regulates Wnt/beta-catenin signaling by translocating DVL into the nucleus (PubMed:25805136). Reduces virus replication, probably by binding the interferon stimulated response element (ISRE) to promote antiviral gene expression (PubMed:25852164). [UniProtKB/Swiss-Prot Function]