

Product datasheet for SC303325

H2AC20 (NM_003517) Human Untagged Clone

Product data:

Product Type: Expression Plasmids

Product Name: H2AC20 (NM_003517) Human Untagged Clone

Tag: Tag Free Symbol: H2AC20

Synonyms: H2A; H2A-GL101; H2A/q; H2AFQ; HIST2H2AC

Mammalian Cell Neo

Selection:

Neomycin

Vector:pCMV6-Entry (PS100001)E. coli Selection:Kanamycin (25 ug/mL)

Fully Sequenced ORF: >SC303325 representing NM_003517.

Blue=Insert sequence Red=Cloning site Green=Tag(s)

GATCCGGTACCGAGGAGATCTGCCGCCGCGATCGCC

ATGTCTGGTCGTGGCAAACAAGGAGGCCAAGGCCCAAGGCCAAGTCGCGCTCGTCCCGCGCTGGC
CTCCAGTTCCCGGTAGGGCGAGTGCACCGCTTGCTGCGCAAAAGGCAACTACGCGGAGCGGGTGGGGGGCC
GGCGCCCCGTCTACATGGCGGCGGTCCTCGAGTACCTGACCGCCGAGATCCTGGAGCTGGCGGCAAC
GCGGCTCGGGACAACAAGAAGAAGACGCGCATCATCCCTCGTCACCTCCAGCTGGCCATCCGCAACGACGAG
GAACTGAACAAGCTGCTGGGCAAAGTCACCATCGCCCAGGGCGGCGTTTTGCCTAACATCCAGGCCGTT
CTGTTACCAAAGAAAACCGAAAAGCCACAAAGCCAAAAGCAAATAA

ACGCGTACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGAT

TACAAGGATGACGACGATAAGGTTTAAACGGCCGGC

Restriction Sites: Sgfl-Mlul


OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Plasmid Map:



ACCN: NM_003517

Insert Size: 390 bp

OTI Disclaimer: Our molecular clone sequence data has been matched to the reference identifier above as a

point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative

RNA splicing form or single nucleotide polymorphism (SNP).

OTI Annotation: This TrueClone is provided through our Custom Cloning Process that includes sub-cloning

into OriGene's pCMV6 vector and full sequencing to provide a non-variant match to the expected reference without frameshifts, and is delivered as lyophilized plasmid DNA.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).



Reconstitution Method:

- 1. Centrifuge at 5,000xg for 5min.
- 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
- 3. Close the tube and incubate for 10 minutes at room temperature.
- 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

RefSeq: <u>NM 003517.2</u>

 RefSeq Size:
 437 bp

 RefSeq ORF:
 390 bp

 Locus ID:
 8338

 UniProt ID:
 Q16777

 Cytogenetics:
 1q21.2

Protein Pathways: Systemic lupus erythematosus

MW: 14 kDa

Gene Summary: Histones are basic nuclear proteins that are responsible for the nucleosome structure of the

chromosomal fiber in eukaryotes. Two molecules of each of the four core histones (H2A, H2B, H3, and H4) form an octamer, around which approximately 146 bp of DNA is wrapped in repeating units, called nucleosomes. The linker histone, H1, interacts with linker DNA between nucleosomes and functions in the compaction of chromatin into higher order structures. This gene is intronless and encodes a replication-dependent histone that is a

member of the histone H2A family. [provided by RefSeq, Aug 2015]