

Product datasheet for **SC303572**

Macrophage Inflammatory Protein 3 (CCL23) (NM_005064) Human Untagged Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	Macrophage Inflammatory Protein 3 (CCL23) (NM_005064) Human Untagged Clone
Tag:	Tag Free
Symbol:	CCL23
Synonyms:	CK-BETA-8; Ckb-8; Ckb-8-1; CKb8; hmrp-2a; MIP-3; MIP3; MPIF-1; SCYA23
Vector:	<u>pCMV6 series</u>
Fully Sequenced ORF:	>NCBI ORF sequence for NM_005064, the custom clone sequence may differ by one or more nucleotides ATGAAGGTCTCCGTGGCTGCCCTCTCCTGCCTCATGCTTGTTACTGCCCTTGGATCCCAG GCCCGGTACAAAAGATGCAGAGACAGAGTTCATGATGTCAAAGCTTCCATTGGAAAAT CCAGTACTTCTGGACATGCTCTGGAGGAGAAAGATTGGTCCTCAGATGACCCTTCTCAT GCTGCAGGATCCATGCTACTAGTGTGCTGACTGCTGCATCTCTACACCCACGAAGCATC CCGTGTTCACTCTGGAGAGTTACTTTGAAACGAACAGCGAGTGTCCAAGCCGGGTGTC ATCTTCTCACCAAGAAGGGGCGACGTTTCTGTGCCAACCCAGTGATAAGCAAGTTCAG GTTTGCCTGAGAATGCTGAAGCTGGACACACGGATCAAGACCAGGAAGAATTGA
Restriction Sites:	Please inquire
ACCN:	NM_005064
OTI Disclaimer:	Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).
OTI Annotation:	This TrueClone is provided through our Custom Cloning Process that includes sub-cloning into OriGene's pCMV6 vector and full sequencing to provide a non-variant match to the expected reference without frameshifts, and is delivered as lyophilized plasmid DNA.
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).



[View online »](#)

Reconstitution Method:	<ol style="list-style-type: none">1. Centrifuge at 5,000xg for 5min.2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.3. Close the tube and incubate for 10 minutes at room temperature.4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
RefSeq:	NM_005064.3 , NP_005055.2
RefSeq Size:	641 bp
RefSeq ORF:	414 bp
Locus ID:	6368
UniProt ID:	P55773
Cytogenetics:	17q12
Protein Families:	Druggable Genome, Secreted Protein
Protein Pathways:	Chemokine signaling pathway, Cytokine-cytokine receptor interaction
Gene Summary:	<p>This gene is one of several chemokine genes clustered on the q-arm of chromosome 17. Chemokines form a superfamily of secreted proteins involved in immunoregulatory and inflammatory processes. The superfamily is divided into four subfamilies based on the arrangement of the N-terminal cysteine residues of the mature peptide. This chemokine, a member of the CC subfamily, displays chemotactic activity on resting T lymphocytes and monocytes, lower activity on neutrophils and no activity on activated T lymphocytes. The protein is also a strong suppressor of colony formation by a multipotential hematopoietic progenitor cell line. In addition, the product of this gene is a potent agonist of the chemokine (C-C motif) receptor 1. Alternative splicing results in multiple transcript variants that encode different isoforms. [provided by RefSeq, Jul 2013]</p> <p>Transcript Variant: This variant (CKbeta8-1) represents the longer transcript and encodes the longer isoform (CKbeta8-1).</p>