

Product datasheet for SC303739

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Septin 2 (SEPT2) (NM_006155) Human Untagged Clone

Product data:

Product Type: Expression Plasmids

Product Name: Septin 2 (SEPT2) (NM_006155) Human Untagged Clone

Tag: Tag Free Symbol: SEPTIN2

Synonyms: DIFF6; hNedd5; NEDD-5; NEDD5; Pnutl3; SEPT2

Vector: pCMV6 series

Fully Sequenced ORF: >NCBI ORF sequence for NM_006155, the custom clone sequence may differ by one or more

nucleotides

ATGTCTAAGCAACAGCCAACTCAGTTTATAAATCCAGAAACACCTGGCTATGTTGGATTT GCAAACCTCCCCAATCAAGTTCACCGAAAATCAGTGAAAAAAGGTTTTGAGTTCACACTG ATGGTGGTCGGTGAATCAGGTCTAGGAAAATCGACTCTCATAAACAGCCTATTCCTAACT GATCTGTACCCAGAAAGAGTCATACCTGGAGCAGCAGAAAAAATTGAAAGAACTGTCCAG ATTGAGGCTTCAACTGTTGAAATTGAAGAGCGAGGGGTCAAGCTACGCCTGACAGTGGTA GATACCCCTGGCTATGGTGACGCTATCAACTGCAGAGATTGTTTTAAGACAATTATCTCC TATATTGATGAGCAATTTGAGAGGTACCTGCATGACGAGAGCGGCTTGAACAGGCGGCAC ATCATTGATAATAGGGTGCATTGTTGCTTTTACTTTATTTCACCTTTTGGACATGGACTT AAGCCCTTAGATGTGGCGTTTATGAAGGCAATACACAACAAGGTGAATATTGTGCCTGTC ATTGCAAAAGCTGACACTCTCACCCTGAAGGAACGGGAGCGGCTGAAGAAAAGGATTCTG GATGAAATTGAAGAACATAACATCAAAATCTATCACTTACCTGATGCAGAATCAGATGAA GATGAAGATTTTAAAGAGCAGACTAGACTTCTCAAGGCTAGCATCCCATTCTCTGTGGTT GGATCCAATCAGTTGATTGAAGCCAAAGGAAAGAAGGTCAGAGGCCGCCTCTACCCCTGG GGTGTTGTGGAAGTGGAGAACCCAGAGCACAATGACTTTCTGAAGCTGAGAACCATGCTC ATCACCCACATGCAGGATCTCCAGGAGGTGACCCAGGACCTTCATTATGAAAACTTCCGT TCTGAGAGACTCAAGAGAGGCGGCAGGAAAGTGGAGAATGAGGACATGAATAAAGACCAG ATCTTGCTGGAAAAAGAAGCTGAGCTCCGCCGCATGCAAGAGATGATTGCAAGGATGCAG GCGCAGATGCAGATGCAGGGCGGGGGGTGGCGGGGCTCTCGGGCACCAC

GTGTAA

Restriction Sites: Please inquire **ACCN:** NM_006155

OTI Disclaimer: Our molecular clone sequence data has been matched to the reference identifier above as a

point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative

RNA splicing form or single nucleotide polymorphism (SNP).





OTI Annotation:

This TrueClone is provided through our Custom Cloning Process that includes sub-cloning into OriGene's pCMV6 vector and full sequencing to provide a non-variant match to the expected reference without frameshifts, and is delivered as lyophilized plasmid DNA.

Components:

The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method:

- 1. Centrifuge at 5,000xg for 5min.
- 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
- 3. Close the tube and incubate for 10 minutes at room temperature.
- 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

RefSeq: <u>NM 006155.1</u>, <u>NP 006146.1</u>

 RefSeq Size:
 3497 bp

 RefSeq ORF:
 1086 bp

 Locus ID:
 4735

 UniProt ID:
 Q15019

 Cytogenetics:
 2q37.3

Gene Summary:

Filament-forming cytoskeletal GTPase. Forms a filamentous structure with SEPTIN12, SEPTIN6, SEPTIN2 and probably SEPTIN4 at the sperm annulus which is required for the structural integrity and motility of the sperm tail during postmeiotic differentiation (PubMed:25588830). Required for normal organization of the actin cytoskeleton. Plays a role in the biogenesis of polarized columnar-shaped epithelium by maintaining polyglutamylated microtubules, thus facilitating efficient vesicle transport, and by impeding MAP4 binding to tubulin. Required for the progression through mitosis. Forms a scaffold at the midplane of the mitotic splindle required to maintain CENPE localization at kinetochores and consequently chromosome congression. During anaphase, may be required for chromosome segregation and spindle elongation. Plays a role in ciliogenesis and collective cell movements. In cilia, required for the integrity of the diffusion barrier at the base of the primary cilium that prevents diffusion of transmembrane proteins between the cilia and plasma membranes: probably acts by regulating the assembly of the tectonic-like complex (also named B9 complex) by localizing TMEM231 protein. May play a role in the internalization of 2 intracellular microbial pathogens, Listeria monocytogenes and Shigella flexneri.[UniProtKB/Swiss-Prot Function] Transcript Variant: This variant (2) differs in the 5' UTR compared to variant 1. Variants 1-4, 8-13, and 16-18 encode the same isoform (a).