

Product datasheet for SC304017

OR10C1 (NM_013941) Human Untagged Clone

Product data:

Product Type: Expression Plasmids

Product Name: OR10C1 (NM_013941) Human Untagged Clone

Tag: Tag Free Symbol: OR10C1

Synonyms: hs6M1-17; OR6-31; OR10C1P; OR10C2

Mammalian Cell

Selection:

Neomycin

Vector:pCMV6-Entry (PS100001)E. coli Selection:Kanamycin (25 ug/mL)

Fully Sequenced ORF: >SC304017 representing NM_013941.

Blue=Insert sequence Red=Cloning site Green=Tag(s)

GATCCGGTACCGAGGAGATCTGCCGCCGCGATCGCC

ACGCGTACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGAT

TACAAGGATGACGACGATAAGGTTTAAACGGCCGGC

Restriction Sites: Sgfl-Mlul

ACCN: NM_013941

Insert Size: 939 bp



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



OR10C1 (NM_013941) Human Untagged Clone - SC304017

OTI Disclaimer: Our molecular clone sequence data has been matched to the reference identifier above as a

point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative

RNA splicing form or single nucleotide polymorphism (SNP).

OTI Annotation: This TrueClone is provided through our Custom Cloning Process that includes sub-cloning

into OriGene's pCMV6 vector and full sequencing to provide a non-variant match to the expected reference without frameshifts, and is delivered as lyophilized plasmid DNA.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method: 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

RefSeq: <u>NM 013941.3</u>

 RefSeq Size:
 1039 bp

 RefSeq ORF:
 939 bp

 Locus ID:
 442194

 UniProt ID:
 Q96KK4

Cytogenetics:

Protein Families: Transmembrane

Protein Pathways: Olfactory transduction

6p22.1

MW: 34.4 kDa

Gene Summary: Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal

response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms. This olfactory receptor gene is a segregating pseudogene, where some individuals have an allele that encodes a functional

olfactory receptor, while other individuals have an allele encoding a protein that is predicted

to be non-functional. [provided by RefSeq, Jul 2015]