

Product datasheet for SC304120

KIR2DS5 (NM_014513) Human Untagged Clone

Product data:


Product Type:	Expression Plasmids
Product Name:	KIR2DS5 (NM_014513) Human Untagged Clone
Tag:	Tag Free
Symbol:	KIR2DS5
Synonyms:	CD158G; NKAT9
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)
Fully Sequenced ORF:	>SC304120 representing NM_014513. Blue=Insert sequence Red=Cloning site Green=Tag(s)

```
GCTCGTTT TAGTGAACCGTCAGAATTTTGT AATACGACTCACTATAGGGCGCCGGGAATTCGTCGACTG
GATCCGGTACCGAGGAGATCTGCCGCC CGCATCGCC
ATGTCGCTCATGGTCATCAGCATGGCGTGTGTTGCGTTCTTCTTGCTGCAGGGGGCCTGGCCACATGAG
GGATTCCGCAGAAAACCTTCCCTCCTGGCCACCCAGGTCCTGGTCAAATCAGAAGAGACAGTCATC
CTGCAATGTTGGTCAGATGTCATGTTTGAGCACTTCTTCTGCACAGAGAGGGGACGTTTAAACCACACT
TTGCGCCTCATTGGAGAGCACATTGATGGGGTCTCCAAGGGCAACTTCTCCATCGGTCGCATGACACAA
GACCTGGCAGGGACCTACAGATGCTACGGTTCTGTTACTCACTCCCCTATCAGTTGTCAGCGCCCACT
GACCCTCTGGACATCGTGATCACAGGTCTATATGAGAAACCTTCTCTCAGCCAGCCGGGCCCCACG
GTTCTGGCAGGAGAGAGCGGTGACCTTGTCTGCAGCTCCCGGAGCTCCTATGACATGTACCATCTATCC
AGGGAAGGGGAGGCCATGAACGTAGGCTCCCTGCAGGGCCCAAGGTCAACAGAACATTCAGGCCGAC
TTTCTCTGGACCCTGCCACCCAGGAGGGACCTACAGATGCTTCGGCTCTTCCGCTGACTCTCCATAC
GAGTGGTCAAAGTCAAAGTACCCACTGCTTGTCTGTACAGGAACTCTTCAAATAGTTGGCCTTCA
CCCCTGAACCAAGTCCGAAACCGGTAAACCCAGACCTACACGTTCTGATTGGGACCTCAGTGGTC
AACTCCCTTTCACCATCCTCCTCTTCTTCTCCTTCATCGCTGGTGTCCAACAAAAAATGCATCT
GTAATGGACCAAGGGCCTGCGGGAACAGAACAGTGAACAGGGAGGATTCTGATGAACAGGACCATCAG
GAGGTGCATACGCATAA
ACGCGTACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGAT
TACAAGGATGACGACGATAAGGTTTAAACGGCCGGC
```

Restriction Sites: SgfI-MluI



[View online »](#)

Plasmid Map:


ACCN: NM_014513

Insert Size: 915 bp

OTI Disclaimer: Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).

OTI Annotation: This TrueClone is provided through our Custom Cloning Process that includes sub-cloning into OriGene's pCMV6 vector and full sequencing to provide a non-variant match to the expected reference without frameshifts, and is delivered as lyophilized plasmid DNA.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method:

1. Centrifuge at 5,000xg for 5min.
2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
3. Close the tube and incubate for 10 minutes at room temperature.
4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

RefSeq: [NM_014513.2](#)

RefSeq Size: 1602 bp

RefSeq ORF: 915 bp

Locus ID:	3810
UniProt ID:	Q14953
Cytogenetics:	19q13.4
Protein Families:	Transmembrane
Protein Pathways:	Antigen processing and presentation, Natural killer cell mediated cytotoxicity
MW:	33.7 kDa
Gene Summary:	<p>Killer cell immunoglobulin-like receptors (KIRs) are transmembrane glycoproteins expressed by natural killer cells and subsets of T cells. The KIR genes are polymorphic and highly homologous and they are found in a cluster on chromosome 19q13.4 within the 1 Mb leukocyte receptor complex (LRC). The gene content of the KIR gene cluster varies among haplotypes, although several "framework" genes are found in all haplotypes (KIR3DL3, KIR3DP1, KIR3DL4, KIR3DL2). The KIR proteins are classified by the number of extracellular immunoglobulin domains (2D or 3D) and by whether they have a long (L) or short (S) cytoplasmic domain. KIR proteins with the long cytoplasmic domain transduce inhibitory signals upon ligand binding via an immune tyrosine-based inhibitory motif (ITIM), while KIR proteins with the short cytoplasmic domain lack the ITIM motif and instead associate with the TYRO protein tyrosine kinase binding protein to transduce activating signals. The ligands for several KIR proteins are subsets of HLA class I molecules; thus, KIR proteins are thought to play an important role in regulation of the immune response. [provided by RefSeq, Jul 2008]</p>