

Product datasheet for SC305804

WFDC3 (NM 080614) Human Untagged Clone

Product data:

Product Type: Expression Plasmids

Product Name: WFDC3 (NM_080614) Human Untagged Clone

Tag: Tag Free Symbol: WFDC3

Synonyms: dJ447F3.3; WAP14

Mammalian Cell

Selection:

None

Vector: pCMV6-XL5

E. coli Selection: Ampicillin (100 ug/mL)

Fully Sequenced ORF: >OriGene sequence for NM_080614 edited

Restriction Sites: Please inquire ACCN: NM 080614

Insert Size: 700 bp

OTI Disclaimer: Our molecular clone sequence data has been matched to the reference identifier above as a

point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative

RNA splicing form or single nucleotide polymorphism (SNP).

OTI Annotation: The ORF of this clone has been fully sequenced and found to contain 2 SNPs compared with

NM 080614.1.



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com

WFDC3 (NM_080614) Human Untagged Clone - SC305804

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method: 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

RefSeq: <u>NM 080614.1, NP 542181.1</u>

 RefSeq Size:
 1000 bp

 RefSeq ORF:
 696 bp

 Locus ID:
 140686

 UniProt ID:
 Q8IUB2

 Cytogenetics:
 20q13.12

Protein Families: Secreted Protein

Gene Summary: This gene encodes a member of the WAP-type four-disulfide core (WFDC) domain family. The

WFDC domain, or WAP signature motif, contains eight cysteines forming four disulfide bonds

at the core of the protein, and functions as a protease inhibitor. The encoded protein

contains four WFDC domains. Most WFDC genes are localized to chromosome 20q12-q13 in

two clusters: centromeric and telomeric. This gene belongs to the telomeric cluster.

Alternatively spliced transcript variants have been observed but their full-length nature has

not been determined. [provided by RefSeq, Jul 2008]