

## Product datasheet for **SC306576**

### SLC22A6 (NM\_153277) Human Untagged Clone

#### Product data:

Product Type:	Expression Plasmids
Product Name:	SLC22A6 (NM_153277) Human Untagged Clone
Tag:	Tag Free
Symbol:	SLC22A6
Synonyms:	HOAT1; OAT1; PAHT; ROAT1
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)
Restriction Sites:	Sgfl-Mlul
ACCN:	NM_153277
Insert Size:	1521 bp
OTI Disclaimer:	Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).
OTI Annotation:	This TrueClone is provided through our Custom Cloning Process that includes sub-cloning into OriGene's pCMV6 vector and full sequencing to provide a non-variant match to the expected reference without frameshifts, and is delivered as lyophilized plasmid DNA.
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	<ol style="list-style-type: none"><li>1. Centrifuge at 5,000xg for 5min.</li><li>2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.</li><li>3. Close the tube and incubate for 10 minutes at room temperature.</li><li>4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.</li><li>5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.</li></ol>
RefSeq:	<a href="#">NM_153277.2</a>



[View online »](#)

RefSeq Size: 2077 bp

RefSeq ORF: 1521 bp

Locus ID: 9356

UniProt ID: [Q4U2R8](#)

Cytogenetics: 11q12.3

Protein Families: Transmembrane

MW: 55.9 kDa

**Gene Summary:** The protein encoded by this gene is involved in the sodium-dependent transport and excretion of organic anions, some of which are potentially toxic. The encoded protein is an integral membrane protein and may be localized to the basolateral membrane. Four transcript variants encoding four different isoforms have been found for this gene. [provided by RefSeq, Jul 2008]

Transcript Variant: This variant (3, also called OAT1-3) lacks two alternate in-frame segments compared to variant 1, resulting in a shorter isoform (c) compared to isoform a.