

## **Product datasheet for SC307018**

## IL5RA (NM\_175727) Human Untagged Clone

## **Product data:**

**Product Type:** Expression Plasmids

Product Name: IL5RA (NM\_175727) Human Untagged Clone

Tag: Tag Free Symbol: IL5RA

Synonyms: CD125; CDw125; HSIL5R3; IL5R

Mammalian Cell No

Selection:

Neomycin

Vector:pCMV6-Entry (PS100001)E. coli Selection:Kanamycin (25 ug/mL)

Fully Sequenced ORF: >SC307018 representing NM\_175727.

Blue=Insert sequence Red=Cloning site Green=Tag(s)

GATCCGGTACCGAGGAGATCTGCCGCCGCGATCGCC

**ACGCGTACGCCGCCCC**GAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGAT

TACAAGGATGACGACGATAAGGTTTAAACGGCCGGC

**Restriction Sites:** Sgfl-Mlul

**ACCN:** NM 175727

**Insert Size:** 1002 bp



**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



## IL5RA (NM\_175727) Human Untagged Clone - SC307018

**OTI Disclaimer:** Our molecular clone sequence data has been matched to the reference identifier above as a

point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative

RNA splicing form or single nucleotide polymorphism (SNP).

OTI Annotation: This TrueClone is provided through our Custom Cloning Process that includes sub-cloning

into OriGene's pCMV6 vector and full sequencing to provide a non-variant match to the expected reference without frameshifts, and is delivered as lyophilized plasmid DNA

**Components:** The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

**Reconstitution Method:** 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

**RefSeq:** <u>NM 175727.1</u>

RefSeq Size: 1983 bp
RefSeq ORF: 1002 bp
Locus ID: 3568

UniProt ID: Q01344

Cytogenetics: 3p26.2

**Protein Families:** Druggable Genome, Transmembrane

**Protein Pathways:** Cytokine-cytokine receptor interaction, Hematopoietic cell lineage, Jak-STAT signaling pathway

**MW:** 37.7 kDa

**Gene Summary:** The protein encoded by this gene is an interleukin 5 specific subunit of a heterodimeric

cytokine receptor. The receptor is comprised of a ligand specific alpha subunit and a signal transducing beta subunit shared by the receptors for interleukin 3 (IL3), colony stimulating factor 2 (CSF2/GM-CSF), and interleukin 5 (IL5). The binding of this protein to IL5 depends on the beta subunit. The beta subunit is activated by the ligand binding, and is required for the biological activities of IL5. This protein has been found to interact with syndecan binding protein (syntenin), which is required for IL5 mediated activation of the transcription factor SOX4. Several alternatively spliced transcript variants encoding four distinct isoforms have

been reported. [provided by RefSeq, Jul 2011]

Transcript Variant: This variant (5) lacks an internal segment within the 5' UTR, and differs in the 3' end region, when compared to variant 1. The resulting isoform (2) has a shorter and

distinct C-terminus, as compared to isoform 1.