

Product datasheet for SC307273

IL24 (NM 181339) Human Untagged Clone

Product data:

Product Type: Expression Plasmids

Product Name: IL24 (NM_181339) Human Untagged Clone

Tag: Tag Free Symbol: IL24

Synonyms: C49A; FISP; IL-24; IL10B; mda-7; MDA7; Mob-5; ST16

Mammalian Cell None

Selection:

Vector: pCMV6-XL5

E. coli Selection: Ampicillin (100 ug/mL)

Fully Sequenced ORF: >OriGene sequence for NM_181339 edited

Restriction Sites: Please inquire **ACCN:** NM_181339

Insert Size: 600 bp

OTI Disclaimer: Our molecular clone sequence data has been matched to the reference identifier above as a

point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative

RNA splicing form or single nucleotide polymorphism (SNP).

OTI Annotation: The ORF of this clone has been fully sequenced and found to be a perfect match to

NM 181339.1.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Reconstitution Method:

- 1. Centrifuge at 5,000xg for 5min.
- 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
- 3. Close the tube and incubate for 10 minutes at room temperature.
- 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

RefSeq: <u>NM 181339.1</u>, <u>NP 851936.1</u>

RefSeq Size: 1633 bp
RefSeq ORF: 147 bp
Locus ID: 11009
Cytogenetics: 1q32.1

Protein Families: Druggable Genome, Secreted Protein

Protein Pathways: Cytokine-cytokine receptor interaction, Jak-STAT signaling pathway

Gene Summary: This gene encodes a member of the IL10 family of cytokines. It was identified as a gene

induced during terminal differentiation in melanoma cells. The protein encoded by this gene can induce apoptosis selectively in various cancer cells. Overexpression of this gene leads to elevated expression of several GADD family genes, which correlates with the induction of apoptosis. The phosphorylation of mitogen-activated protein kinase 14 (MAPK7/P38), and heat shock 27kDa protein 1 (HSPB2/HSP27) are found to be induced by this gene in melanoma cells, but not in normal immortal melanocytes. Alternatively spliced transcript variants encoding distinct isoforms have been reported. [provided by RefSeq, Jul 2008] Transcript Variant: This variant (2) lacks an internal segment, as compared to variant 1. The

resulting shorter isoform (2) is identical to the C-terminal region of isoform 1.