

## Product datasheet for **SC307374**

### UBE2C (NM\_181800) Human Untagged Clone

#### Product data:

Product Type:	Expression Plasmids
Product Name:	UBE2C (NM_181800) Human Untagged Clone
Tag:	Tag Free
Symbol:	UBE2C
Synonyms:	dj447F3.2; UBCH10
Vector:	<u>pCMV6 series</u>
Fully Sequenced ORF:	>NCBI ORF sequence for NM_181800, the custom clone sequence may differ by one or more nucleotides ATGGCTTCCCAAACCGCGACCCAGCCGCCACTAGCGTCGCCGCCCGCTAAAGGAGCT GAGCCGAGCGGGGGCGCCGCCGGGTCCGGTGGGCAAAGGCTACAGCAGGAGCTGATG ACCCTCATGGTATATGAAGACCTGAGGTATAAGCTCTCGCTAGAGTTCCCCAGTGGCTAC CCTTACAATGCGCCACAGTGAAGTTCCTCAGCCCTGCTATCACCCAACGTGGACACC CAGGGTAAACATATGCCTGGACATCCTGAAGGAAAAGTGGTCTGCCCTGTATGATGTCAGG ACCATTCTGCTCTCCATCCAGAGCCTTCTAGGAGAACCCAACATTGATAGTCCCTTGAAC ACACATGTGCCGAGCTCTGGAAAACCCACAGCTTTTAAGAAGTACCTGCAAGAAACC TACTCAAAGCAGGTCACCAGCCAGGAGCCCTGA
Restriction Sites:	Please inquire
ACCN:	NM_181800
OTI Disclaimer:	Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).
OTI Annotation:	This TrueClone is provided through our Custom Cloning Process that includes sub-cloning into OriGene's pCMV6 vector and full sequencing to provide a non-variant match to the expected reference without frameshifts, and is delivered as lyophilized plasmid DNA.
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).



[View online »](#)

**Reconstitution Method:**

1. Centrifuge at 5,000xg for 5min.
2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
3. Close the tube and incubate for 10 minutes at room temperature.
4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

**RefSeq:** [NM\\_181800.1](#), [NP\\_861516.1](#)

**RefSeq Size:** 744 bp

**RefSeq ORF:** 453 bp

**Locus ID:** 11065

**UniProt ID:** [O00762](#)

**Cytogenetics:** 20q13.12

**Protein Families:** Druggable Genome

**Protein Pathways:** Ubiquitin mediated proteolysis

**Gene Summary:** The modification of proteins with ubiquitin is an important cellular mechanism for targeting abnormal or short-lived proteins for degradation. Ubiquitination involves at least three classes of enzymes: ubiquitin-activating enzymes, ubiquitin-conjugating enzymes, and ubiquitin-protein ligases. This gene encodes a member of the E2 ubiquitin-conjugating enzyme family. The encoded protein is required for the destruction of mitotic cyclins and for cell cycle progression, and may be involved in cancer progression. Multiple transcript variants encoding different isoforms have been found for this gene. Pseudogenes of this gene have been defined on chromosomes 4, 14, 15, 18, and 19. [provided by RefSeq, Aug 2013]  
Transcript Variant: This variant (3) lacks an in-frame exon in the central coding region, compared to variant 1. The encoded isoform (3) is shorter, compared to isoform 1.