

Product datasheet for **SC307498**

CREM (NM_182724) Human Untagged Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	CREM (NM_182724) Human Untagged Clone
Tag:	Tag Free
Symbol:	CREM
Synonyms:	CREM-2; hCREM-2; ICER
Vector:	<u>pCMV6 series</u>
Fully Sequenced ORF:	>NCBI ORF sequence for NM_182724, the custom clone sequence may differ by one or more nucleotides ATGAACAGAACTCAGGAGTTGTCTGGCCAGCTTAGTGCTGCCACTGGTGACATGCCAACT TACCAGATCCGAGCTCCTACTGCTGCTTTGCCACAGGGAGTGGTGATGGCTGCATCGCCC GGAAGTTTGCACAGTCCCAGCAGCTGGCAGAAGAAGCAACACGCAAACGAGAGCTGAGG CTAATGAAAAACAGGGAAGCTGCCAAAGAATGTCGACGTCGAAAGAAAGAATATGTAAAA TGTCTGGAGAGCCGAGTTGCAGTGCTGGAAGTCCAGAACAAGAAGCTTATAGAGGAACTT GAAACCTTGAAAGACATTTGTTCTCCCAAACAGATTACTAG
Restriction Sites:	Please inquire
ACCN:	NM_182724
OTI Disclaimer:	Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).
OTI Annotation:	This TrueClone is provided through our Custom Cloning Process that includes sub-cloning into OriGene's pCMV6 vector and full sequencing to provide a non-variant match to the expected reference without frameshifts, and is delivered as lyophilized plasmid DNA.
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).



[View online »](#)

Reconstitution Method:

1. Centrifuge at 5,000xg for 5min.
2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
3. Close the tube and incubate for 10 minutes at room temperature.
4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

RefSeq: [NM_182724.1](#), [NP_874393.1](#)

RefSeq Size: 1663 bp

RefSeq ORF: 342 bp

Locus ID: 1390

UniProt ID: [Q03060](#)

Cytogenetics: 10p11.21

Protein Families: Druggable Genome, Transcription Factors

Gene Summary: This gene encodes a bZIP transcription factor that binds to the cAMP responsive element found in many viral and cellular promoters. It is an important component of cAMP-mediated signal transduction during the spermatogenic cycle, as well as other complex processes. Alternative promoter and translation initiation site usage allows this gene to exert spatial and temporal specificity to cAMP responsiveness. Multiple alternatively spliced transcript variants encoding several different isoforms have been found for this gene, with some of them functioning as activators and some as repressors of transcription. [provided by RefSeq, Jul 2008]

Transcript Variant: This variant (11) uses an alternative downstream promoter, and differs in the 5' UTR and 5' coding region, compared to variant 1. This results in a shorter isoform (11, also known as k) with a distinct N-terminus, compared to isoform 1. This isoform represents one of the early repressor ICER isoforms. Sequence Note: This RefSeq record was created from transcript and genomic sequence data to make the sequence consistent with the reference genome assembly. The genomic coordinates used for the transcript record were based on transcript alignments.