

Product datasheet for **SC307841**

SYT10 (NM_198992) Human Untagged Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	SYT10 (NM_198992) Human Untagged Clone
Tag:	Tag Free
Symbol:	SYT10
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)



[View online »](#)

Fully Sequenced ORF: >SC307841 representing NM_198992.
 Blue=Insert sequence Red=Cloning site Green=Tag(s)

```

GCTCGTTTAGTGAACCGTCAGAATTTTGTAAACGACTCACTATAGGGCGGCCGGGAATTCGTCGACTG
GATCCGGTACCGAGGAGATCTGCCGCCGCGATCGCC
ATGAGTTTCCACAAGGAGGACGGAGTGAACAGTCTGTGCCAGAAGGCTCTGCACATCGTCACCGAGCTG
TGCTTCGCCGGCCAGGTGGAGTGGGAGAGTGTCTGGGCATCTTCCCTCGGGACAGGGGCAGCCAGGGC
GGAAGCAGCACAGATATTTCAAGTCAGCCTGTTAGCTGTCGTTGTACAGCTTTTGTGGACTGGCCTTGTG
GTTGTCTCACTTTTGTCTTCTGGAAGCTGTGTTGGCCATGCTGGAAGCAAACTGTGACTTCCAAC
ATCACTACGCTTCCACAGAGCATTTCAAGTGCTCTACTGAAGTTTTTGAGACTGAAGAGAAAAAGAA
ATTAAGGAAAAAGAAAGCCAGCCGTAAGAGCTATTGAGCCTGCAATAAAATCAGCCACACTTCCCTT
GACATCCAGCAGAGTCCAACTGCTTTAAAGAACATTTAATTAACATGCACGTGTGCAAGACAA
ATTACTGAGCCTACGTCATCAACCCGCCACAGTTCCTTCCGAAGACACCTGCCGAGGCAATGCAGGTT
TCCAGTGTGATTTTAGCATGGGCACAGAACCTGTTTACAACGAGGAGAAACAACACAGCATTGGG
AGGATAAAGCCAGAACTCTACAAACAGAAATCAGTTGACTCTGAGGGCAACCAAAACGAAGATGTCAA
ATCTGTGGGAACTTAACCTTACCCTCCAGTATGATTATGAAATGAAGTCTAGTTGTTAAATATC
AAAGCTTTAGATCTCCCTGCTAAAGACTTCACAGGAAGTCTGACCCCTTATGTGAAGATGTATCTCTT
CCAGATAGGAAAAAGAAATTTAGACCCCGGTGCACAGAAAGACTTTAAATCCTCTATTTGATGAACT
TTTCAATTTCTGTAGCATATGATCAACTAAGCAACCGAAACTACATTTAGTGTGTATGATTTTGAC
AGATTTTCTAGACATGACATGATTGGGGAAGTGATTCTTGATAATTTGTTGAAGTCTCTGATCTCTCC
AGGGAAGCCACAGTATGGAAAGATTTCACTGTGCTACCAAGAAAGTATAGACCTGGGTGAAATCATG
TTTTCCCTTTGTACCTACCGACGGCTGGGCGTATGACATTGACAGTCATTAAGTGCAGAAATCTGAAG
GCGATGGATATTACTGGCTCATCAGATCCTTATGTCAAAGTGTCCCTGATGTGTGAAGTCAAGATTA
AAAAAGAGGAAAAACAACCTACAAAGAAAAACACTTAAACCCTGTGTACAATGAGGCCATTATTTTGAC
ATCCCTCCAGAGAACGTGGACCGGTGAGCCTCTCCATTGCGGTCATGGATTACGATAGGGTAGGACAC
AATGAGGTGATAGGAGTGTGCAGAACAGGACTGGATGCTGAGGGTCTTGGGCGAGACCACTGGAATGAA
ATGCTGGCCTATCATCGAAACCAATAACGCACTGGCACCCATTGCTGGAGTTACCTGGCCGGGCGAGC
AGTTTTGATAGTCAAGGATCCTGCCCTTCTCTAAACCACCTTCCACACCAAA
ACGCGTACGCGGCCGCTCGAGCAGAACTCATCTCAGAGAGGATCTGGCAGCAATGATATCTGGAT
TACAAGGATGACGACGATAAGGTTTAAACGGCCGGC
  
```

Restriction Sites: SgfI-MluI

ACCN: NM_198992

Insert Size: 1572 bp

OTI Disclaimer: Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).

OTI Annotation: This TrueClone is provided through our Custom Cloning Process that includes sub-cloning into OriGene's pCMV6 vector and full sequencing to provide a non-variant match to the expected reference without frameshifts, and is delivered as lyophilized plasmid DNA.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method:	1. Centrifuge at 5,000xg for 5min. 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA. 3. Close the tube and incubate for 10 minutes at room temperature. 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom. 5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
RefSeq:	<u>NM_198992.3</u>
RefSeq Size:	3286 bp
RefSeq ORF:	1572 bp
Locus ID:	341359
UniProt ID:	<u>Q6XYQ8</u>
Cytogenetics:	12p11.1
Protein Families:	Secreted Protein, Transmembrane
MW:	59.1 kDa
Gene Summary:	Ca(2+) sensor specifically required for the Ca(2+)-dependent exocytosis of secretory vesicles containing IGF1 in neurons of the olfactory bulb. Exocytosis of IGF1 is required for sensory perception of smell. Not involved in Ca(2+)-dependent synaptic vesicle exocytosis (By similarity). Acts through Ca(2+) and phospholipid binding to the C2 domain: Ca(2+) induces binding of the C2-domains to phospholipid membranes and to assembled SNARE-complexes; both actions contribute to triggering exocytosis (By similarity).[UniProtKB/Swiss-Prot Function]