

Product datasheet for **SC308080**

TCF7 (NM_201634) Human Untagged Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	TCF7 (NM_201634) Human Untagged Clone
Tag:	Tag Free
Symbol:	TCF7
Synonyms:	TCF-1
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)
Fully Sequenced ORF:	>SC308080 representing NM_201634. Blue=Insert sequence Red=Cloning site Green=Tag(s)

GCTCGTTTAGTGAACCGTCAGAATTTTGAATACGACTCACTATAGGGCGCCGGGAATTCGTCGACTG
 GATCCGGTACCGAGGAGATCTGCCGCC**GCGATCGCC**
 ATGTACAAAGAGACCGTCTACTCCGCCTTCAATCTGCTCATGCATTACCCACCCCTCGGGAGCAGGG
 CAGCACCCCGAGCCGAGCCCCGCTGCACAAGGCCAATCAGCCCCCGACGGTGTCCCCAACTCTCT
 CTCTACGAACATTTCAACAGCCACATCCACCCCTGCACCTGCGGACATCAGCCAGAAGCAAGTTCAC
 AGGCCTCTGCAGACCCCTGACCTCTCTGGCTTCTACTCCCTGACCTCAGGCAGCATGGGGCAGCTCCCC
 CACACTGTGAGCTGGTTACCCACCCATCCTTGATGCTAGGTTCTGGTGTACCTGGTCACCCAGCAGCC
 ATCCCCACCCGCCATTGTGCCCCCTCAGGGAAGCAGGAGCTGCAGCCCTTCGACCGCAACCTGAAG
 ACACAAGCAGAGTCCAAGGCAGAGAAGGAGGCCAAGAAGCCAACCATCAAGAAGCCCTCAATGCCTTC
 ATGCTGTACATGAAGGAGATGAGAGCCAAGGTCATTGCAGAGTGACACTTAAGGAGAGCGCTGCCATC
 AACCAGATCCTGGGCCGAGGTGGCACGCGCTGTCGCGAGAAGAGCAGGCCAAGTACTATGAGCTGGCC
 CGCAAGGAGAGGCAGCTGCACATGCAGCTATACCCAGGCTGGTCAGCGCGGGACAACCTACGGGAAGAAG
 AAGAGGCGGTGAGGGAAAAGCACAAGAATCCACCACAGACCCTGGCTCGCCTAAGAAATGCCGTGCT
 CGCTTTGGCCTCAACCAGCAGACGGATTGGTGTGGTCCGTGCAGTA**TAA**
ACGCGTACGCGGCCGCTCGAGCAGAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGAT
 TACAAGGATGACGACGATAAGGTTTAAACGGCCGGC

Restriction Sites:	SgfI-MluI
ACCN:	NM_201634
Insert Size:	807 bp


[View online »](#)

OTI Disclaimer:	Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).
OTI Annotation:	This TrueClone is provided through our Custom Cloning Process that includes sub-cloning into OriGene's pCMV6 vector and full sequencing to provide a non-variant match to the expected reference without frameshifts, and is delivered as lyophilized plasmid DNA.
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	<ol style="list-style-type: none"> 1. Centrifuge at 5,000xg for 5min. 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA. 3. Close the tube and incubate for 10 minutes at room temperature. 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom. 5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
RefSeq:	<u>NM_201634.3</u>
RefSeq Size:	2935 bp
RefSeq ORF:	807 bp
Locus ID:	6932
UniProt ID:	<u>P36402</u>
Cytogenetics:	5q31.1
Protein Families:	Druggable Genome, ES Cell Differentiation/IPS, Transcription Factors
Protein Pathways:	Acute myeloid leukemia, Adherens junction, Arrhythmogenic right ventricular cardiomyopathy (ARVC), Basal cell carcinoma, Colorectal cancer, Endometrial cancer, Melanogenesis, Pathways in cancer, Prostate cancer, Thyroid cancer, Wnt signaling pathway
MW:	30.3 kDa

Gene Summary:

This gene encodes a member of the T-cell factor/lymphoid enhancer-binding factor family of high mobility group (HMG) box transcriptional activators. This gene is expressed predominantly in T-cells and plays a critical role in natural killer cell and innate lymphoid cell development. The encoded protein forms a complex with beta-catenin and activates transcription through a Wnt/beta-catenin signaling pathway. Mice with a knockout of this gene are viable and fertile, but display a block in T-lymphocyte differentiation. Alternative splicing results in multiple transcript variants. Naturally-occurring isoforms lacking the N-terminal beta-catenin interaction domain may act as dominant negative regulators of Wnt signaling. [provided by RefSeq, Oct 2016]

Transcript Variant: This variant (4, also known as C), differs in the 5' UTR, has multiple coding region differences, uses a downstream start codon, and differs in the 3' UTR, compared to variant 1. The resulting isoform (4) is shorter at the N-terminus and has a distinct C-terminus, compared to isoform 1. Sequence Note: This RefSeq record was created from transcript and genomic sequence data to make the sequence consistent with the reference genome assembly. The genomic coordinates used for the transcript record were based on transcript alignments.