

Product datasheet for **SC308088**

splicing factor 1 (SF1) (NM_201995) Human Untagged Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	splicing factor 1 (SF1) (NM_201995) Human Untagged Clone
Tag:	Tag Free
Symbol:	SF1
Synonyms:	BBP; D11S636; MBBP; ZCCHC25; ZFM1; ZNF162
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)



[View online »](#)

Fully Sequenced ORF: >SC308088 representing NM_201995.
 Blue=Insert sequence Red=Cloning site Green=Tag(s)

```

GCTCGTTTAGTGAACCGTCAGAATTTTGTAAACGACTCACTATAGGGCGGCCGGGAATTCGTGACTG
GATCCGGTACCGAGGAGATCTGCCGCCCGGATCGCC
ATGGCGACCGGAGCGAACGCCACGCCGTTGGACTTCCCAAGTAAAGAGCGGAAGAGGAGCCGCTGGAAC
CAAGACACAATGGAACAGAAGACAGTGATTCCAGGAATGCCTACAGTTATCCCCCTGGACTTACTCGA
GAACAAGAAAGAGCTTATATAGTGCAACTGCAGATAGAAGACCTGACTCGTAAACTGCGCACAGGAGAC
CTGGGCATCCCCCTAACCTGAGGACAGGTCCCTTCCCTGAGCCATCTACAATAGCGAGGGGAAG
CGGCTTAACACCCGAGAGTTCGCCACCCGCAAAAAGCTGGAAGAGGAGCGGCACAACCTCATCACAGAG
ATGTTGCACTCAATCCGGATTTCAAGCCACCTGCAGATTACAAACCTCCAGCAACACGTGTGAGTGAT
AAAGTCATGATCCACAAGATGAGTACCAGAAATCAACTTTGTGGGGCTGCTCATCGGGCCAGAGGG
AACACCTGAAGAACATAGAGAAGGAGTGAATGCCAAGATTATGATCCGGGGAAAGGGTCTGTGAAA
GAAGGGAAGTTGGCGCAAAGATGGCCAGATGTTGCCAGGAGAAGATGAGCCACTTCATGCCCTGGTT
ACTGCCAATAACAATGGAGAACGTCAAAAAGGCAGTGAACAGATAAGAAACATCCTGAAGCAGGGTATC
GAGACTCCAGAGGACCAGAATGATCTACGGAAGATGCAGCTTCGGGAGTTGGCTCGCTTAAATGGGACC
CTTCGGGAAGACGATAACAGGATCTTAAGACCCTGGCAGAGCTCAGAGACCCGCAGCATTACCAACACC
ACAGTGTGTACCAAGTGTGGAGGGGCTGGCCACATTGCTTCAGACTGTAAATCCAAAGGCTGGTGAT
CCTCAGTCAGCTCAGGATAAAGCACGGATGGATAAAGAATATTTGCCCTCATGGCTGAAGTGGGTGAA
GCACCTGTCCCAGCATCTGTGGGCTCCACCTCTGGGCTGCCACCACACCCCTGGCCAGCGCACCTCGT
CCTGCTGCTCCGCCAACACCCACCTCCACCGTCTCTCATGTCTACCACCCAGAGCCGCCACCCCTGG
ATGAATCTGGCCCTCAGAGAGTGGCCCTACCACGGCATGCATGGAGTGGTCTCTGGTGGGCCGGGA
GGTGGCCCCACAGCTTCCCACACCCATTACCCAGCCTGACAGGTGGGCATGGTGGACATCCCATGCAG
CACAAACCAATGGACCCACCCCTTGGATGCAGCCACCACCACCCAGATGAACCAGGGCCCCCAC
CCTCCTGGGACCATGGCCCTCCTCCAATGGATCAGTACCTGGGAAGTACGCCTGTGGGCTCTGGGGTC
TATCGCTGCATCAAGGAAAAGGTATGATGCCGCCACCACCTATGGGCATGATGCCGCCGCCGCCGCCG
CCTCCCAGTGGGACGCCCCACCCCTCCCTCTGGTCTCTTCCCCATGGCAACAACAGCAGCAGCAG
CCTCCGCCACCCCTCCGCCAGCAGCAGTATGGCTTCCAGTACCCCTTGGCATGGCAGCAAAATACG
ACGACTACCACACGAGCGCTGGCACAGGGTCCATCCCGCATGGCAACAGCAGCAGGGCGGTGCCGCA
GCTTCTCCAGGAGCCCTCAGATGCAAGGCAACCCACTATGGTGGCCCTGCCCCCGGGTCCAGCCG
CCTCTGCCGCTGGGGCCCTCCCTCCGCCCTCCACCGCTGGTTCGGCCGCATGATGATCCCT
CCCGCGCGGGGATGGCCCGAGCCATGAGAGTAGGACTTCCGCGCCATTGGTGACCCTTCCAGGC
AGACAGCCTCAGCAACGCCCTGGTGGACAGGATGGTTCGGCAAAGCAGCCTGA
ACGCGTACGCGGCCGCTCGAGCAGAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGAT
TACAAGGATGACGACGATAAGGTTTAAACGGCCGGC
  
```

Restriction Sites: SgfI-MluI

Plasmid Map: □

ACCN: NM_201995

Insert Size: 1917 bp

OTI Disclaimer: Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).

OTI Annotation: This TrueClone is provided through our Custom Cloning Process that includes sub-cloning into OriGene's pCMV6 vector and full sequencing to provide a non-variant match to the expected reference without frameshifts, and is delivered as lyophilized plasmid DNA.

Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	<ol style="list-style-type: none">1. Centrifuge at 5,000xg for 5min.2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.3. Close the tube and incubate for 10 minutes at room temperature.4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
RefSeq:	NM_201995.2
RefSeq Size:	3154 bp
RefSeq ORF:	1917 bp
Locus ID:	7536
UniProt ID:	Q15637
Cytogenetics:	11q13.1
Protein Families:	Transcription Factors
MW:	68.6 kDa
Gene Summary:	<p>This gene encodes a nuclear pre-mRNA splicing factor. The encoded protein specifically recognizes the intron branch point sequence at the 3' splice site, together with the large subunit of U2 auxiliary factor (U2AF), and is required for the early stages of spliceosome assembly. It also plays a role in nuclear pre-mRNA retention and transcriptional repression. The encoded protein contains an N-terminal U2AF ligand motif, a central hnRNP K homology motif and quaking 2 region which bind a key branch-site adenosine within the branch point sequence, a zinc knuckles domain, and a C-terminal proline-rich domain. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Oct 2016]</p> <p>Transcript Variant: This variant (2) lacks a segment in the 3' UTR and coding region which results in a frameshift, compared to variant 1. It encodes isoform 2, which is shorter and has a distinct C-terminus, compared to isoform 1.</p>