

Product datasheet for SC308285

QK1 (QKI) (NM_206853) Human Untagged Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	QK1 (QKI) (NM_206853) Human Untagged Clone
Tag:	Tag Free
Symbol:	QKI
Synonyms:	Hqk; hqkl; QK; QK1; QK3
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)
Fully Sequenced ORF:	>SC308285 representing NM_206853. Blue=Insert sequence Red=Cloning site Green=Tag(s)

```
GCTCGTTTGTAGTGAACCGTCAGAATTTTGTAAACGACTCACTATAGGGCCGCCGGGAATTCGTGACTG
GATCCGGTACCGAGGAGATCTGCCGCCGCGATCGCC
ATGGTCGGGAAATGAAACGAAGGAGAAGCCGAAGCCACCCAGATTACCTGATGCAGCTGATGAAC
GACAAGAAGCTCATGAGCAGCCTGCCAACTTCTGCGGGATCTTCAACCACCTCGAGCGGCTGCTGGAC
GAAGAAATTAGCAGAGTACGGAAGACATGTACAATGACACATTAATGGCAGTACAGAGAAAAGGAGT
GCAGAATTGCCTGATGCTGTGGGACCTATTGTTTCAGTTACAAGAGAAAACCTTATGTGCCTGTAAAAGAA
TACCCAGATTTTAAATTTGTTGGGAGAATCCTTGGACCTAGAGGACTTACAGCCAACAACCTGAAGCA
GAAACCGGATGTAAAATCATGGTCCGAGGCAAAGGCTCAATGAGGGATAAAAAAAGGAGGAGCAAAT
AGAGGCAAGCCCAATTGGGAGCATCTAAATGAAGATTTACATGTACTAATCACTGTGGAAGATGCTCAG
AACAGAGCAGAAAATCAAATTGAAGAGAGCAGTTGAAGAAGTGAAGAAAATTATTGGTACCTGCAGCAGAA
GGAGAAGCAGCCTGAAGAAGATGCAGCTGATGGAGCTTGCGATTCTGAATGGCACCTACAGAGATGCC
AACATTAATCACCAGCCCTTGCCTTTTCTCTTGACGCAACAGCCAGGCTGCTCCAAGGATCATTACT
GGGCTGCGCCGGTTCTCCACCAGCTGCCCTGCGTACTCCTACGCCAGCTGGCCCTACCATAATGCCT
TTGATCAGACAAATACAGACCGCTGTATGCCAAACGGAACCTCCTACCCAACCTGCTGCAATAGTTCTT
CCAGGCCCCGAAGCTGGTTAACTATAACCCCTATGAGTACCCCTACACATTGGCACCAGCTACATCA
ATCCTTGAGTATCCTATTGAACCTAGTGGTGTATTAGGTATGGCTTTCCCAACGAAAGGCTAA
ACGCGTACGCGGCCGCTCGAGCAGAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGAT
TACAAGGATGACGACGATAAGGTTTAAACGCCCGGC
```

Restriction Sites:	Sgfl-Mlul
Plasmid Map:	<input type="checkbox"/>
ACCN:	NM_206853



[View online »](#)

Insert Size:	960 bp
OTI Disclaimer:	Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).
OTI Annotation:	This TrueClone is provided through our Custom Cloning Process that includes sub-cloning into OriGene's pCMV6 vector and full sequencing to provide a non-variant match to the expected reference without frameshifts, and is delivered as lyophilized plasmid DNA.
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	<ol style="list-style-type: none">1. Centrifuge at 5,000xg for 5min.2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.3. Close the tube and incubate for 10 minutes at room temperature.4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
RefSeq:	NM_206853.2
RefSeq Size:	14152 bp
RefSeq ORF:	960 bp
Locus ID:	9444
UniProt ID:	Q96PU8
Cytogenetics:	6q26
MW:	35.1 kDa
Gene Summary:	<p>The protein encoded by this gene is an RNA-binding protein that regulates pre-mRNA splicing, export of mRNAs from the nucleus, protein translation, and mRNA stability. The encoded protein is involved in myelination and oligodendrocyte differentiation and may play a role in schizophrenia. Multiple transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Jul 2014]</p> <p>Transcript Variant: This variant (2) differs in the 3' UTR and coding region compared to variant 1, which results in a frameshift. The resulting isoform (2; also known as HQK-6 and QKI-6) is shorter and has a distinct C-terminus compared to isoform 1. Isoforms 2 and 4 are the same length but have different C-termini. Sequence Note: This RefSeq record was created from transcript and genomic sequence data to make the sequence consistent with the reference genome assembly. The genomic coordinates used for the transcript record were based on transcript alignments.</p>