

Product datasheet for SC308707

INCA1 (NM 213726) Human Untagged Clone

Product data:

Product Type: Expression Plasmids

Product Name: INCA1 (NM_213726) Human Untagged Clone

Tag: Tag Free
Symbol: INCA1
Synonyms: HSD45

Mammalian Cell

Selection:

Neomycin

Vector:pCMV6-Entry (PS100001)E. coli Selection:Kanamycin (25 ug/mL)

Fully Sequenced ORF: >SC308707 representing NM_213726.

Blue=Insert sequence Red=Cloning site Green=Tag(s)

GATCCGGTACCGAGGAGATCTGCCGCCGCGATCGCC

ACGCGTACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGAT

TACAAGGATGACGACGATAAGGTTTAAACGGCCGGC

Restriction Sites: Sgfl-Mlul ACCN: NM_213726

Insert Size: 666 bp

OTI Disclaimer: Our molecular clone sequence data has been matched to the reference identifier above as a

point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative

RNA splicing form or single nucleotide polymorphism (SNP).



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com ORIGENE

OTI Annotation: This TrueClone is provided through our Custom Cloning Process that includes sub-cloning

into OriGene's pCMV6 vector and full sequencing to provide a non-variant match to the expected reference without frameshifts, and is delivered as lyophilized plasmid DNA.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method: 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

17p13.2

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

RefSeq: <u>NM 213726.2</u>

 RefSeq Size:
 1404 bp

 RefSeq ORF:
 666 bp

 Locus ID:
 388324

 UniProt ID:
 Q0VD86

MW: 25.2 kDa

Cytogenetics:

Gene Summary: Binds to CDK2-bound cyclins and inhibits the kinase activity of CDK2; binding to cyclins is

critical for its function as CDK inhibitor (PubMed:21540187). Inhibits cell growth and cell proliferation and may play a role in cell cycle control (By similarity). Required for ING5-mediated regulation of S-phase progression, enhancement of Fas-induced apoptosis and

inhibition of cell growth (By similarity).[UniProtKB/Swiss-Prot Function]

Transcript Variant: This variant (3) differs in the 5' UTR and uses an alternate in-frame splice site in the coding region, compared to variant 1. This results in a shorter protein (isoform 2),

compared to isoform 1. Both variants 3 and 4 encode the same isoform (2).