

Product datasheet for **SC308801**

FGFR1 (NM_023107) Human Untagged Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	FGFR1 (NM_023107) Human Untagged Clone
Tag:	Tag Free
Symbol:	FGFR1
Synonyms:	BFGFR; CD331; CEK; FGFBR; FLG; FLJ99988; FLT2; HBGFR; KAL2; N-SAM; OGD
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)
Fully Sequenced ORF:	>SC308801 representing NM_023107. Blue=Insert sequence Red=Cloning site Green=Tag(s)

```
GCTCGTTTGTAGTGAACCGTCAGAATTTGTAAACGACTACTATAGGGCGCCGGGAATTCGTGACTG
GATCCGGTACCGAGGAGATCTGCCGCCGCGATCGCC
ATGTGGAGCTGGAAGTGCCTCCTCTTCTGGGCTGTGCTGGTACACGCCACACTCTGCACCCTAGGCCG
TCCCCGACCTTGCTGAACAAGATGCTCTCCCCTCCTCGGAGGATGATGATGATGATGACTCCTCT
TCAGAGGAGAAAGAAACAGATAACACCAAACAAACCGTATGCCCGTAGCTCCATATTGGACATCCCCA
GAAAAGATGAAAAGAAATTGCATGCAGTGCCGGCTGCCAAGACAGTGAAGTTCAAATGCCCTTCCAGT
GGGACCCAAACCCACACTGCGCTGGTTGAAAAATGGCAAAGAATTCAAACCTGACCACAGAATTGGA
GGCTACAAGTCCGTTATGCCACCTGGAGCATATAATGGACTCTGTGGTGGCCCTGACAAGGGCAAC
TACACCTGCATTGTGGAGAATGAGTACGGCAGCATCAACCACACATACCAGCTGGATGTCGTGGAGCGG
TCCCCTACCCGGCCCATCCTGCAAGCAGGGTTGCCCGCCAAACAAACAGTGGCCCTGGGTAGCAACGTG
GAGTTCATGTGTAAGGTGTACAGTGACCCGAGCCGCACATCCAGTGGCTAAAGCACATCGAGGTGAAT
GGGAGCAAGATTGGCCAGACAACCTGCCTTATGTCCAGATCTTGAAGTAATCATGGCACCAGTCTTC
GTGGGCCAGTCTACTGGGAAGGAGACACTGTCTCGGGGCTCAAGTTCTGTGGCAGGCTCAGTTGC
CCCCGAATGGGATCATTCTCACGCTTCAGGCACACACTCCATCTCAGTAGGGATCTAGCCACATCC
CCCAGGACTAGTAACAGAGGTACAAAGTGGAGGTGAGCTGGGAACAGAGGGCTGCAGGGATGGTGGT
GCTGGTCTGTAA
ACGCGTACGCGCCGCTCGAGCAGAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGAT
TACAAGGATGACGACGATAAGGTTTAAACGGCCGGC
```

Restriction Sites:	Sgfl-MluI
Plasmid Map:	<input type="checkbox"/>
ACCN:	NM_023107



[View online »](#)

Insert Size:	909 bp
OTI Disclaimer:	Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).
OTI Annotation:	This TrueClone is provided through our Custom Cloning Process that includes sub-cloning into OriGene's pCMV6 vector and full sequencing to provide a non-variant match to the expected reference without frameshifts, and is delivered as lyophilized plasmid DNA.
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	<ol style="list-style-type: none"> 1. Centrifuge at 5,000xg for 5min. 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA. 3. Close the tube and incubate for 10 minutes at room temperature. 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom. 5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
RefSeq:	<u>NM_023107.2</u>
RefSeq Size:	2807 bp
RefSeq ORF:	909 bp
Locus ID:	2260
Cytogenetics:	8p11.23
Domains:	ig, IGc2, IG
Protein Families:	Druggable Genome, Protein Kinase, Transmembrane
Protein Pathways:	Adherens junction, MAPK signaling pathway, Melanoma, Pathways in cancer, Prostate cancer, Regulation of actin cytoskeleton
MW:	33.4 kDa

Gene Summary:

The protein encoded by this gene is a member of the fibroblast growth factor receptor (FGFR) family, where amino acid sequence is highly conserved between members and throughout evolution. FGFR family members differ from one another in their ligand affinities and tissue distribution. A full-length representative protein consists of an extracellular region, composed of three immunoglobulin-like domains, a single hydrophobic membrane-spanning segment and a cytoplasmic tyrosine kinase domain. The extracellular portion of the protein interacts with fibroblast growth factors, setting in motion a cascade of downstream signals, ultimately influencing mitogenesis and differentiation. This particular family member binds both acidic and basic fibroblast growth factors and is involved in limb induction. Mutations in this gene have been associated with Pfeiffer syndrome, Jackson-Weiss syndrome, Antley-Bixler syndrome, osteoglophonic dysplasia, and autosomal dominant Kallmann syndrome 2. Chromosomal aberrations involving this gene are associated with stem cell myeloproliferative disorder and stem cell leukemia lymphoma syndrome. Alternatively spliced variants which encode different protein isoforms have been described; however, not all variants have been fully characterized. [provided by RefSeq, Jul 2008]

Transcript Variant: This variant (5) differs in the 3' UTR and lacks multiple exons in the 3' coding region, which results in an early stop codon, compared to variant 1. This variant encodes isoform 5, also known as isoform H4, which is much shorter, has distinct C-terminus, and lacks the transmembrane domain, compared to isoform 1.