

## **Product datasheet for SC308888**

## OGG1 (NM 016819) Human Untagged Clone

**Product data:** 

**Product Type:** Expression Plasmids

**Product Name:** OGG1 (NM\_016819) Human Untagged Clone

Tag: Tag Free Symbol: OGG1

Synonyms: HMMH; HOGG1; MUTM; OGH1

Mammalian Cell

Selection:

Neomycin

Vector:pCMV6-Entry (PS100001)E. coli Selection:Kanamycin (25 ug/mL)

Fully Sequenced ORF: >SC308888 representing NM\_016819.

Blue=Insert sequence Red=Cloning site Green=Tag(s)

GATCCGGTACCGAGGAGATCTGCCGCCGCGATCGCC

**ACGCGTACGCGGCCGCTC**GAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGAT

TACAAGGATGACGACGATAAGGTTTAAACGGCCGGC

**Restriction Sites:** Sgfl-Mlul

Plasmid Map:

**ACCN:** NM 016819



**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com

## OGG1 (NM\_016819) Human Untagged Clone - SC308888

**Insert Size:** 975 bp

**OTI Disclaimer:** Our molecular clone sequence data has been matched to the reference identifier above as a

point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative

RNA splicing form or single nucleotide polymorphism (SNP).

OTI Annotation: This TrueClone is provided through our Custom Cloning Process that includes sub-cloning

into OriGene's pCMV6 vector and full sequencing to provide a non-variant match to the expected reference without frameshifts, and is delivered as lyophilized plasmid DNA.

**Components:** The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

**Reconstitution Method:** 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

**RefSeq:** <u>NM 016819.3</u>

 RefSeq Size:
 1896 bp

 RefSeq ORF:
 975 bp

 Locus ID:
 4968

 UniProt ID:
 015527

Cytogenetics: 3p25.3

Protein Families: Druggable Genome
Protein Pathways: Base excision repair

MW: 36.4 kDa

**Gene Summary:** This gene encodes the enzyme responsible for the excision of 8-oxoguanine, a mutagenic

base byproduct which occurs as a result of exposure to reactive oxygen. The action of this enzyme includes lyase activity for chain cleavage. Alternative splicing of the C-terminal region of this gene classifies splice variants into two major groups, type 1 and type 2, depending on the last exon of the sequence. Type 1 alternative splice variants end with exon 7 and type 2 end with exon 8. All variants share the N-terminal region in common, which contains a mitochondrial targeting signal that is essential for mitochondrial localization. Many alternative splice variants for this gene have been described, but the full-length nature for

every variant has not been determined. [provided by RefSeq, Aug 2008]

Transcript Variant: Transcript variant 1b contains a 244 bp insertion between exons 6 and 7,

as compared to the predominant transcript variant 1a.