

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for SC309015

Lymphotactin (XCL1) (NM_002995) Human Untagged Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	Lymphotactin (XCL1) (NM_002995) Human Untagged Clone
Tag:	Tag Free
Symbol:	Lymphotactin
Synonyms:	ATAC; LPTN; LTN; SCM-1; SCM-1a; SCM1; SCM1A; SCYC1
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)
Fully Sequenced ORF:	>NCBI ORF sequence for NM_002995, the custom clone sequence may differ by one or more nucleotides
	ATGAGACTTCTCATCCTGGCCCTCCTTGGCATCTGCTCTCTCACTGCATACATTGTGGAAGGTGTAGGGA GTGAAGTCTCAGATAAGAGGACCTGTGTGAGCCTCACTACCCAGCGACTGCCGGTTAGCAGAATCAAGAC CTACACCATCACGGAAGGCTCCTTGAGAGCAGTAATTTTTATTACCAAACGTGGCCTAAAAGTCTGTGCT GATCCACAAGCCACATGGGTGAGAGACGTGGTCAGGAGCATGGACAGGAAATCCAACACCAGAAATAACA TGATCCAGACCAAGCCAACAGGAACCCAGCAATCGACCAATACAGCTGTGACTCTGACTGGCTAG
Restriction Sites:	Please inquire
ACCN:	



View online »

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

	nphotactin (XCL1) (NM_002995) Human Untagged Clone – SC309015
OTI Disclaimer:	Due to the inherent nature of this plasmid, standard methods to replicate additional amounts of DNA in E. coli are highly likely to result in mutations and/or rearrangements. Therefore, OriGene does not guarantee the capability to replicate this plasmid DNA. Additional amounts of DNA can be purchased from OriGene with batch-specific, full-sequence verification at a reduced cost. Please contact our customer care team at <u>custsupport@origene.com</u> or by calling 301.340.3188 option 3 for pricing and delivery.
	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <u>More info</u>
OTI Annotation:	This TrueClone is provided through our Custom Cloning Process that includes sub-cloning into OriGene's pCMV6 vector and full sequencing to provide a non-variant match to the expected reference without frameshifts, and is delivered as lyophilized plasmid DNA.
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Meth	 od: 1. Centrifuge at 5,000xg for 5min. 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA. 3. Close the tube and incubate for 10 minutes at room temperature. 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom. 5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
RefSeq:	<u>NM 002995.1, NP 002986.1</u>
RefSeq Size:	1222 bp
RefSeq ORF:	345 bp
Locus ID:	6375
UniProt ID:	<u>P47992</u>
Cytogenetics:	1q24.2
Domains:	IL8
Protein Families:	Druggable Genome, Secreted Protein
Protein Pathways:	Chemokine signaling pathway, Cytokine-cytokine receptor interaction

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

	Lymphotactin (XCL1) (NM_002995) Human Untagged Clone – SC309015
Gene Summary:	This antimicrobial gene encodes a member of the chemokine superfamily. Chemokines function in inflammatory and immunological responses, inducing leukocyte migration and activation. The encoded protein is a member of the C-chemokine subfamily, retaining only two of four cysteines conserved in other chemokines, and is thought to be specifically chemotactic for T cells. This gene and a closely related family member are located on the long

arm of chromosome 1. [provided by RefSeq, Sep 2014]

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US