

Product datasheet for **SC309753**

BAX (NM_004324) Human Untagged Clone

Product data:

Product Type: Expression Plasmids
Product Name: BAX (NM_004324) Human Untagged Clone
Tag: Tag Free
Symbol: BAX
Synonyms: BCL2L4
Mammalian Cell Selection: None
Vector: pCMV6-XL5
E. coli Selection: Ampicillin (100 ug/mL)

Fully Sequenced ORF: >OriGene ORF sequence for NM_004324 edited
 GGGCGGCCGCGAATTCGCCCTTGGAGCGGCGGTGATGGACGGTCCGGGGAGCAGCCCAG
 AGGCGGGGGGCCACCAGCTCTGAGCAGATCATGAAGACAGGGGCCCTTTGCTTCAGGG
 TTTTCATCCAGGATCGAGCAGGGCGAATGGGGGGGAGGCACCCGAGCTGGCCCTGGACCC
 GGTGCCTCAGGATGCGTCCACCAAGAAGCTGAGCGAGTGTCTCAAGCGCATCGGGGACGA
 ACTGGACAGTAACATGGAGCTGCAGAGGATGATTGCCGCCGTGGACACAGACTCCCCCG
 AGAGGTCTTTTTCCGAGTGGCAGCTGACATGTTTTCTGACGGCAACTTCAACTGGGGCCG
 GGTTGTCGCCCTTTTCTACTTTGCCAGCAAAGTGGTGCTCAAGGCCCTGTGCACCAAGGT
 GCCGGAAGTATCAGAACCATCATGGGCTGGACATTGGACTTCTCCGGGAGCGGCTGTT
 GGGCTGGATCCAAGACCAGGGTGGTTGGGTGAGACTCCTCAAGCCTCCTACCCCCACCA
 CCGCGCCCTCACCACCGCCCTGCCACCGTCCCTGCCCGCCACTCCTCTGGGACC
 CTGGGCCTTCTGGAGCAGGTACAGTGGTCCCTCTCCCATCTTCAGATCATCAGATGT
 GGTCTATAATGCGTTTTCTTACGTGTCTGATCAATCCCCGATTCATCTACCCTGCTGAC
 CTCCAGTGACCCTGACAAGGGCGAATTCAGATCTGGTACCGATATCAAGCTTGTGCGAC
 TCTAGA



[View online »](#)

5' Read Nucleotide Sequence:	>OriGene 5' read for NM_004324 unedited GGGCGGCAAGTTTTGTATACGACTCACTATAGGGCGGCCGCGATTGCGCCCTTGGAGCGGC GGTGATGGACGGGTCCGGGGAGCAGCCCAGAGGCGGGGGCCACCAGCTCTGAGCATTC ATGAAGCAGGGGCCCTTTTGCTTCAGGGTTTCATCCAGGATCGAGCAGGGCGAATGGGGG GGGAGGCACCCGAGCTGGCCCTGGACCCGGTGCCTCAGGATGCGTCCACCAAGAAGCTGA GCGAGTGTCTCAAGCGCATCGGGGACGAACTGGACAGTAACATGGAGCTGCAGAGGATGA TTGCCCGCGTGGACACAGACTCCCCCGAGAGGTCTTTTTCCGAGTGGCAGCTGACATGT TTTCTGACGGCAACTTCAACTGGGGCCGGTTGTGCGCCCTTTTCTACTTTGCCAGCAAAC TGGTGCTCAAGGCCCTGTGCACCAAGGTGCCGGAAGTATCAGAACCATCATGGGCTGGA CATTGGACTTCTCCGGGAGCGGCTGTTGGGCTGGATCCAAGACCAGGTGGTTGGGTGA GACTCCTCAAGCCTCCTACCCCCACCACCGCCCTCACCACCGCCCTGCCCCACCGT CCCTGCCCCCGCCACTCCTCTGGGACCCTGGGCCTTCTGGAGCAGGTACAGTGGTGCC CTCTCCCCATCTCAGATCATCAGATGTGGTCTATAATGCGTTTTCTTACGTGTCTGAT CAATCCCGATTATCTACCCTGCTGACCTCCCAGTGACCCTGACAAGGGCGAATTCAG ATCTGGTACCGATATCAAGCTTGTGACTCTAGATTGCGGCCGCGGTATAGCTGTTTCC TGAACAGATCCCGGTGGCATCCCTGTGACCCCTCCCAGTGCTCTCTGGC
Restriction Sites:	Please inquire
ACCN:	NM_004324
OTI Disclaimer:	Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).
OTI Annotation:	The open reading frame of this TrueClone was fully sequenced and found to be a perfect match to the protein associated to this reference.
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	<ol style="list-style-type: none"> 1. Centrifuge at 5,000xg for 5min. 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA. 3. Close the tube and incubate for 10 minutes at room temperature. 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom. 5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
RefSeq:	NM_004324.3 , NP_004315.1
RefSeq Size:	891 bp
RefSeq ORF:	657 bp
Locus ID:	581
UniProt ID:	Q07812
Cytogenetics:	19q13.33
Domains:	Bcl-2
Protein Families:	Druggable Genome, Transmembrane

Protein Pathways:	Amyotrophic lateral sclerosis (ALS), Apoptosis, Colorectal cancer, Huntington's disease, Neurotrophin signaling pathway, p53 signaling pathway, Pathways in cancer, Prion diseases
Gene Summary:	<p>The protein encoded by this gene belongs to the BCL2 protein family. BCL2 family members form hetero- or homodimers and act as anti- or pro-apoptotic regulators that are involved in a wide variety of cellular activities. This protein forms a heterodimer with BCL2, and functions as an apoptotic activator. The association and the ratio of BAX to BCL2 also determines survival or death of a cell following an apoptotic stimulus. This protein is reported to interact with, and increase the opening of, the mitochondrial voltage-dependent anion channel (VDAC), which leads to the loss in membrane potential and the release of cytochrome c. The expression of this gene is regulated by the tumor suppressor P53 and has been shown to be involved in P53-mediated apoptosis. Multiple alternatively spliced transcript variants, which encode different isoforms, have been reported for this gene. [provided by RefSeq, Dec 2019]</p> <p>Transcript Variant: This variant (beta) differs in the 3' coding region and 3' UTR, compared to variant 1. The resulting isoform (beta) has a shorter and distinct C-terminus, compared to isoform 1.</p>