

Product datasheet for **SC309942**

ATP2B1 (NM_001682) Human Untagged Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	ATP2B1 (NM_001682) Human Untagged Clone
Tag:	Tag Free
Symbol:	ATP2B1
Synonyms:	PMCA1; PMCA1kb
Mammalian Cell Selection:	None
Vector:	<u>pCMV6-XL5</u>
E. coli Selection:	Ampicillin (100 ug/mL)

Fully Sequenced ORF: >OriGene sequence for NM_001682 edited
GCCACCATGGGCGACATGGCAAACAACACTCAGTTGCTTACAGTGGTGTGAAAACTCTTTG
AAGGAAGCTAATCATGATGGAGACTTTTGAATTACGCTCGCAGAGCTGCGGGCTCTCATG
GAGCTCAGGTCCACAGATGCATTACGAAAAATACAGGAAAGCTATGGAGATGTCTATGGA
ATTTGCACCAAATTGAAAACATCTCCAATGAAGGTTTAAAGTGGAAACCTGCAGATTTA
GAAAGAAGAGAAGCAGTGTGGAAAGAATTTTATACCTCCTAAAAAGCCAAAAACCTTT
CTTCAATTAGTATGGGAAGCATTACAAGATGTCACCTTAAATTATATTAGAAATTGCAGCC
ATAGTATCATTGGGCCTTTCTTTTTATCAGCCTCCAGAAGGGGATAATGCACTTTGTGGA
GAAGTTTCTGTTGGGGAGGAAGAAGGTGAAGGTGAAACTGGTTGGATTGAAGGAGCTGCA
ATCCTCTGTCTGTAGTGTGTGTGGTGTAGTAACAGCTTCAATGACTGGAGTAAGGAA
AAACAGTTTAGAGTTTGCAGAGCCGAATTGAACAAGAACAGAAGTTCAGTGCATCAGG
GGTGGTCAAGTCAATCAGATACCTGTAGCTGACATTACTGTTGGAGATATTGCTCAAGTG
AAATATGGTGATCTTCTCCAGCTGACGGCATACTTATCAAGGCAACGATCTTAAAAATT
GATGAAAGCTCATTGACTGGTGAATCAGATCATGTTAAAAAGTCTTTAGATAAGGATCCC
TTACTTCTATCAGGTAATGTAATGGAAGGCTCTGGAAGAATGGTAGTTACAGCTGTA
GGTGTAATTTCTCAAAGTGAATATCTTTACCTTACTTGGAGCTGGAGGTGAAGAGGAA
GAGAAGAAAGATGAGAAGAAAAAGGAAAAAGAAAAATAAGAAACAAGATGGAGCTATTGAG
AATCGCAACAAAAGCAAAGCCAGGATGGTGCAGCCATGGAAATGCAGCCATTGAAGAGT
GAAGAAGGTGGAGATGGTATGAAAAAGATAAAAAGAAAGCAAATTTGCCAAAAAGGAA
AAATCTGTTTTACAAGGAAACTTACAAAAGTGGCTGTTTCAGATTGGCAAAGCAGGCTCTG
TTGATGTCTGCCATCACAGTTATCATTCTAGTATTATATTTTTGTCATTGACACCTTCTGG
GTTCAGAAAAAGCCATGGCTTGCTGAGTGCACACCAATTTATATAACAATACTTTGTGAAG
TTCTTCATTATTGGAGTTACAGTTTTAGTGGTCGAGTGCAGAGGCTTCCACTTGCA
GTCACGATCTCACTGGCTTATTCAGTCAAAAAATGATGAAAGATAATAACTTAGTAAGG
CATCTGGATGCTTGTGAAACCATGGGAAATGCTACAGCTATTTGTTTCAGATAAACAGGA
ACTTTGACAATGAACAGAATGACAGTCGTTCAAGCTTACATAAATGAAAAACATTATAAA
AAGTTTCTGAACCAGAAGCTATTCCACCAAAATTTTTGTCCTATCTTGAACAGGAATT



[View online »](#)

```
TCTGTGAATTGTGCTTATACATCAAAAATTGCCACCAGAGAAAAGGGTGGATTACCT
CGTCACGTTGGTAATAAACTGAATGTGCCTTGTGGGACTTCTTTTGGATTTAAAACGG
GATTATCAGGATGTTAGAAATGAAATACCAGAAGAAGCACTGTACAAAGTCTACACCTTC
AATTCTGTTAGGAAGTCCATGAGTACTGTCTGAAAAATTCAGATGGAAGTTATCGAATA
TTCAGCAAGGGTGCATCTGAGATAATTCTGAAAAAGTGTTCAAAATCTTGAGTGCTAAT
GGTGAGGCAAAAAGTATTCAGACCAAGGGACCGTGATGATATTGAAAACTGTGATTGAA
CCGATGGCATCAGAAGGCTTGAGAACCATATGTCTTGCAATTCAGAGATTTCCAGCAGGA
GAACCAGAACCAGAGTGGGATAATGAAAATGATATTGTCACCGGCCTTACATGCATTGCT
GTTGTGGGGATTGAAGATCCTGTGAGACCTGAGGTGCCAGATGCAATTA AAAAGTGCAG
AGGGCTGGAATTACTGTGCGGATGGTCACTGGTGATAATTAATACTGCTCGGGCCATT
GCTACCAATGTGGTATTTTACATCCTGGGGAAGATTTTCTGTGCCTAGAAGGTAAGAT
TTTAACAGAAGAATACGAAATGAAAAAGGAGAGATTGAGCAAGAGAGGATAGACAAGATT
TGGCCAAAACCTCGAGTACTTGAAGATCATCTCCTACTGATAAGCATACACTGGTTAAA
GGTATAATTGACAGCACTGTCTCAGACCAACGCCAGGTTGTAGCTGTAAGTGGTATGGT
ACAAATGATGGCCAGCACTAAAGAAAGCAGATGTTGGATTTGCAATGGGTATTGCTGGA
ACTGATGTAGCTAAAGAAGCATCCGATATTATTCTCACAGATGACAACTTTACAAGCATT
GTTAAAGCAGTTATGTGGGGACGAAATGTCTATGACAGCATCTCAAAATTCCTTCAGTTC
CAACTTACTGTTAATGTAGTAGCAGTGATTGTTGCTTTTACGGGCGCCTGCATTACTCAA
GACTCACCGCTTAAGGCTGTGCAGATGTGTGGGTAAACCTCATAATGGATACACTCGCT
TCCCTGGCTCTGGCAACGGAACCCACTGAGTCTCTCTTGCTTCGGAACCTTATGGT
AGAAATAAGCCTCTCATCTCACGTACAATGATGAAGAATTTTGGGTCATGCATTCTAT
CAACTTGTAGTAGTCTTTACACTCTTATTTGCTGGAGAAAAGTTTTTGGACATTGATAGT
GGAAGAAATGCTCCTTTGTCATGCTCCTCCTTCAGAACATTATACTATTGTTTTTAATACC
TTTGTGCTGATGCAACTTTTCAACGAAATAAATGCCCGGAAAATTCATGGTGAAAGAAAT
GTATTCGAAGGAATCTTTAACAATGCCATCTTCTGCACAATTGTTTTAGGCACTTTTGTG
GTACAGATAATAATTGTGCAGTTTGGTGGAAAACCTTTCAGTTGTTTCAAGAACTTCAATA
GAACAGTGGCTATGGTCAATATTCCTAGGAATGGGAACATTACTCTGGGGCCAGCTTATT
TCAACAATCCAAGTCCGTTTAAAAATTCCTCAAAGAAGCTGGTCAAGAACACAAAAG
GAAGAAATACCTGAGGAGGAATTAGCAGAGGATGTTGAAGAGATTGATCACGCTGAAAGG
GAGTTGCCGCGTGCCAAATCTTGTGGTTAGAGGTCTGAACAGAATCCAACACAGATT
CGAGTGGTGAATGCATTTCTGAT
```

Restriction Sites:

Please inquire

ACCN:

NM_001682

Insert Size:

3300 bp

OTI Disclaimer:

Due to the inherent nature of this plasmid, standard methods to replicate additional amounts of DNA in *E. coli* are highly likely to result in mutations and/or rearrangements. Therefore, OriGene does not guarantee the capability to replicate this plasmid DNA. Additional amounts of DNA can be purchased from OriGene with batch-specific, full-sequence verification at a reduced cost. Please contact our customer care team at custsupport@origene.com or by calling 301.340.3188 option 3 for pricing and delivery.

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. [More info](#)

OTI Annotation:	This TrueClone is provided through our Custom Cloning Process that includes sub-cloning into OriGene's pCMV6 vector and full sequencing to provide a non-variant match to the expected reference without frameshifts, and is delivered as lyophilized plasmid DNA.
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	<ol style="list-style-type: none">1. Centrifuge at 5,000xg for 5min.2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.3. Close the tube and incubate for 10 minutes at room temperature.4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
RefSeq:	NM_001682.2 , NP_001673.2
RefSeq Size:	6796 bp
RefSeq ORF:	3663 bp
Locus ID:	490
UniProt ID:	P20020
Cytogenetics:	12q21.33
Domains:	E1-E2_ATPase, Cation_ATPase_N, Hydrolase, Cation_ATPase_C
Protein Families:	Druggable Genome, Transmembrane
Protein Pathways:	Calcium signaling pathway
Gene Summary:	<p>The protein encoded by this gene belongs to the family of P-type primary ion transport ATPases characterized by the formation of an aspartyl phosphate intermediate during the reaction cycle. These enzymes remove bivalent calcium ions from eukaryotic cells against very large concentration gradients and play a critical role in intracellular calcium homeostasis. The mammalian plasma membrane calcium ATPase isoforms are encoded by at least four separate genes and the diversity of these enzymes is further increased by alternative splicing of transcripts. The expression of different isoforms and splice variants is regulated in a developmental, tissue- and cell type-specific manner, suggesting that these pumps are functionally adapted to the physiological needs of particular cells and tissues. This gene encodes the plasma membrane calcium ATPase isoform 1. Alternatively spliced transcript variants encoding different isoforms have been identified. [provided by RefSeq, Jul 2008]</p> <p>Transcript Variant: This variant (2) lacks a segment in the coding region, which leads to a frameshift, compared to variant 1. The resulting isoform (1b) contains a longer and distinct C-terminus compared to isoform 1a. Variants 2, 3, 4, and 5 all encode the same isoform (1b).</p> <p>Sequence Note: The RefSeq transcript and protein were derived from genomic sequence to make the sequence consistent with the reference genome assembly. The genomic coordinates used for the transcript record were based on alignments.</p>