

## Product datasheet for **SC310558**

### p38 (MAPK14) (NM\_139013) Human Untagged Clone

#### Product data:

|                           |   |
|---------------------------|---|
| Product Type:             | Expression Plasmids   |
| Product Name:             | p38 (MAPK14) (NM_139013) Human Untagged Clone   |
| Tag:                      | Tag Free  |
| Symbol:                   | MAPK14  |
| Synonyms:                 | CSBP; CSBP1; CSBP2; CSPB1; EXIP; Mxi2; p38; p38ALPHA; PRKM14; PRKM15; RK; SAPK2A        |
| Mammalian Cell Selection: | Neomycin  |
| Vector:                   | pCMV6-Entry (PS100001)  |
| E. coli Selection:        | Kanamycin (25 ug/mL)  |
| Fully Sequenced ORF:      | >SC310558 representing NM_139013.<br>Blue=Insert sequence Red=Cloning site Green=Tag(s) |


```
GCTCGTTTGTAGTGAACCGTCAGAATTTTGTAAACGACTACTATAGGGCGCCGGGAATTCGTGACTG
GATCCGGTACCGAGGAGATCTGCCGCCGCGATCGCC
ATGTCTCAGGAGAGGCCACGTTCTACCGCAGGAGCTGAACAAGACAATCTGGGAGGTGCCGAGCGT
TACCAGAACCTGTCTCCAGTGGGCTCTGGCGCCTATGGCTCTGTGTGTGCTGCTTTTACACAAAAACG
GGGTTACGTGTGGCAGTGAAGAAGCTCTCCAGACCATTTTCAGTCCATCATTATGCGAAAAGAACCTAC
AGAGAACTGCGGTTACTTAAACATATGAAACATGAAAAATGTGATTGGTCTGTTGGACGTTTTTACACCT
GCAAGGTCTCTGGAGGAATCAATGATGTGTATCTGGTGACCCATCTCATGGGGCAGATCTGAACAAC
ATTGTGAAATGTCAGAAGCTTACAGATGACCATGTTTCAGTTCCTTATCTACCAAATTCCTCGAGGTCTA
AAGTATATACATTCAGCTGACATAATTCACAGGGACCTAAAACCTAGTAATCTAGCTGTGAATGAAGAC
TGTGAGCTGAAGATTCTGGATTTTGGACTGGCTCGGCACACAGATGATGAAATGACAGGCTACGTGGCC
ACTAGGTGGTACAGGGCTCCTGAGATCATGCTGAACTGGATGCATTACAACCAGACAGTTGATATTTGG
TCAGTGGGATGCATAATGGCCGAGCTGTTGACTGGAAGAACATGTTTCTGGTACAGACCATATTGAT
CAGTTGAAGCTCATTTTAAAGACTCGTTGGAACCCAGGGGCTGAGCTTTTGAAGAAAATCTCCTCAGAG
TCTGCAAGAACTATATTCAGTCTTTGACTCAGATGCCGAAGTGAACCTTGGCAATGTATTTATTGGT
GCCAATCCCCTGGTAAGTTGACCATATATCCTCACCTCATGGATATTGAATTGGTTATGATATAA
ACGCGTACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGAT
TACAAGGATGACGACGATAAGGTTTAAACGGCCGCGC
```

Restriction Sites: SgfI-MluI



[View online »](#)

Plasmid Map:



ACCN: NM\_139013

Insert Size: 894 bp

OTI Disclaimer: Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).

OTI Annotation: This TrueClone is provided through our Custom Cloning Process that includes sub-cloning into OriGene's pCMV6 vector and full sequencing to provide a non-variant match to the expected reference without frameshifts, and is delivered as lyophilized plasmid DNA.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method: 

1. Centrifuge at 5,000xg for 5min.
2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
3. Close the tube and incubate for 10 minutes at room temperature.
4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

RefSeq: [NM\\_139013.2](#)

RefSeq Size: 1431 bp

RefSeq ORF: 894 bp

|                          |   |
|--------------------------|---|
| <b>Locus ID:</b>         | 1432  |
| <b>UniProt ID:</b>       | <a href="#">Q16539</a>  |
| <b>Cytogenetics:</b>     | 6p21.31   |
| <b>Domains:</b>          | pkinase, TyrKc, S_TKc   |
| <b>Protein Families:</b> | Druggable Genome, Protein Kinase  |
| <b>Protein Pathways:</b> | Amyotrophic lateral sclerosis (ALS), Epithelial cell signaling in Helicobacter pylori infection, Fc epsilon RI signaling pathway, GnRH signaling pathway, Leukocyte transendothelial migration, MAPK signaling pathway, Neurotrophin signaling pathway, NOD-like receptor signaling pathway, Progesterone-mediated oocyte maturation, RIG-I-like receptor signaling pathway, T cell receptor signaling pathway, Toll-like receptor signaling pathway, VEGF signaling pathway  |
| <b>MW:</b>               | 34.1 kDa  |
| <b>Gene Summary:</b>     | <p>The protein encoded by this gene is a member of the MAP kinase family. MAP kinases act as an integration point for multiple biochemical signals, and are involved in a wide variety of cellular processes such as proliferation, differentiation, transcription regulation and development. This kinase is activated by various environmental stresses and proinflammatory cytokines. The activation requires its phosphorylation by MAP kinase kinases (MKKs), or its autophosphorylation triggered by the interaction of MAP3K7IP1/TAB1 protein with this kinase. The substrates of this kinase include transcription regulator ATF2, MEF2C, and MAX, cell cycle regulator CDC25B, and tumor suppressor p53, which suggest the roles of this kinase in stress related transcription and cell cycle regulation, as well as in genotoxic stress response. Four alternatively spliced transcript variants of this gene encoding distinct isoforms have been reported. [provided by RefSeq, Jul 2008]</p> <p>Transcript Variant: This variant (3) contains a different internal segment within the coding region, and a different 3' coding region as well as a different 3' UTR, when compared to variant 1. It thus encodes an isoform that has a different internal segment, and a distinct C-terminus, as compared to isoform 1.</p> |