

Product datasheet for SC311075

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

ERK1 (MAPK3) (NM_001040056) Human Untagged Clone

Product data:

Product Type: Expression Plasmids

Product Name: ERK1 (MAPK3) (NM_001040056) Human Untagged Clone

Tag: Tag Free Symbol: MAPK3

Synonyms: ERK-1; ERK1; ERT2; HS44KDAP; HUMKER1A; p44-ERK1; p44-MAPK; P44ERK1; P44MAPK; PRKM3

Mammalian Cell

Neomycin

Selection:

Vector:pCMV6-Entry (PS100001)E. coli Selection:Kanamycin (25 ug/mL)

Fully Sequenced ORF: >SC311075 representing NM_001040056.

Blue=Insert sequence Red=Cloning site Green=Tag(s)

GATCCGGTACCGAGGAGATCTGCCGCCGCGATCGCC

GTCCCGGGGGAGGTGGAGATGGTGAAGGGGCAGCCGTTCGACGTGGGCCCGCGCTACACGCAGTTGCAG TACATCGGCGAGGGCGCGTACGGCATGGTCAGCTCGGCCTATGACCACGTGCGCAAGACTCGCGTGGCC ATCAAGAAGATCAGCCCCTTCGAACATCAGACCTACTGCCAGCGCACGCTCCGGGAGATCCAGATCCTG CTGCGCTTCCGCCATGAGAATGTCATCGGCATCCGAGACATTCTGCGGGCGTCCACCCTGGAAGCCATG AGAGATGTCTACATTGTGCAGGACCTGATGGAGACTGACCTGTACAAGTTGCTGAAAAGCCAGCAGCTG AGCAATGACCATATCTGCTACTTCCTCTACCAGATCCTGCGGGGCCTCAAGTACATCCACTCCGCCAAC GTGCTCCACCGAGATCTAAAGCCCTCCAACCTGCTCATCAACACCACCTGCGACCTTAAGATTTGTGAT TTCGGCCTGGCCCGGATTGCCGATCCTGAGCATGACCACACCGGCTTCCTGACGGAGTATGTGGCTACG CGCTGGTACCGGGCCCCAGAGATCATGCTGAACTCCAAGGGCTATACCAAGTCCATCGACATCTGGTCT GTGGGCTGCATTCTGGCTGAGATGCTCTCTAACCGGCCCATCTTCCCTGGCAAGCACTACCTGGATCAG CTCAACCACATTCTGGGCATCCTGGGCTCCCCATCCCAGGAGGACCTGAATTGTATCATCAACATGAAG GCCCGAAACTACCTACAGTCTCTGCCCTCCAAGACCAAGGTGGCTTGGGCCAAGCTTTTCCCCAAGTCA GACTCCAAAGCCCTTGACCTGCTGGACCGGATGTTAACCTTTAACCCCAATAAACGGATCACAGTGGAG GAAGCGCTGGCTCACCCCTACCTGGAGCAGTACTATGACCCGACGGATGAGGTGGGCCAGTCCCCAGCA GCAGTGGGGCTGGGGGCAGGGGAGCAGGGGGGCACGTAG

ACGCGTACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGAT


TACAAGGATGACGACGATAAGGTTTAAACGGCCGGC

Restriction Sites: Sgfl-Mlul





Plasmid Map:



ACCN: NM 001040056

Insert Size: 1074 bp

OTI Disclaimer: Our molecular clone sequence data has been matched to the reference identifier above as a

point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative

RNA splicing form or single nucleotide polymorphism (SNP).

OTI Annotation: This TrueClone is provided through our Custom Cloning Process that includes sub-cloning

into OriGene's pCMV6 vector and full sequencing to provide a non-variant match to the expected reference without frameshifts, and is delivered as lyophilized plasmid DNA.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method: 1. Centrifuge at 5,000xg for 5min.

5595

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

RefSeq: NM 001040056.2

RefSeq Size: 1174 bp RefSeq ORF: 1074 bp

Locus ID:

 UniProt ID:
 P27361

 Cytogenetics:
 16p11.2

Protein Families: Druggable Genome, ES Cell Differentiation/IPS, Protein Kinase

Protein Pathways: Acute myeloid leukemia, Adherens junction, Alzheimer's disease, Axon guidance, B cell

receptor signaling pathway, Bladder cancer, Chemokine signaling pathway, Chronic myeloid leukemia, Colorectal cancer, Dorso-ventral axis formation, Endometrial cancer, ErbB signaling

pathway, Fc epsilon RI signaling pathway, Fc gamma R-mediated phagocytosis, Focal adhesion, Gap junction, Glioma, GnRH signaling pathway, Insulin signaling pathway, Long-

term depression, Long-term potentiation, MAPK signaling pathway, Melanogenesis,

Melanoma, mTOR signaling pathway, Natural killer cell mediated cytotoxicity, Neurotrophin signaling pathway, NOD-like receptor signaling pathway, Non-small cell lung cancer, Oocyte meiosis, Pancreatic cancer, Pathways in cancer, Prion diseases, Progesterone-mediated oocyte maturation, Prostate cancer, Regulation of actin cytoskeleton, Renal cell carcinoma, T cell receptor signaling pathway, TGF-beta signaling pathway, Thyroid cancer, Toll-like receptor signaling pathway, Type II diabetes mellitus, Vascular smooth muscle contraction, VEGF

signaling pathway

MW: 40.1 kDa

Gene Summary: The protein encoded by this gene is a member of the MAP kinase family. MAP kinases, also

known as extracellular signal-regulated kinases (ERKs), act in a signaling cascade that regulates various cellular processes such as proliferation, differentiation, and cell cycle progression in response to a variety of extracellular signals. This kinase is activated by upstream kinases, resulting in its translocation to the nucleus where it phosphorylates nuclear targets. Alternatively spliced transcript variants encoding different protein isoforms

have been described. [provided by RefSeg, Jul 2008]

Transcript Variant: This variant (2, also known as ERK1c) lacks several exons and its 3'-terminal exon extends past a splice site that is used in variant 1. The encoded isoform (2) has a shorter

and distinct C-terminus, compared to isoform 1.