

Product datasheet for SC311257

PTPN20A (NM_001042395) Human Untagged Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	PTPN20A (NM_001042395) Human Untagged Clone
Tag:	Tag Free
Symbol:	PTPN20A
Synonyms:	bA142I17.1; CT126; hPTPN20; PTPN20B
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)
Fully Sequenced ORF:	>SC311257 representing NM_001042395. Blue=Insert sequence Red=Cloning site Green=Tag(s)


```
GCTCGTTTAGTGAACCGTCAGAATTTTGTAAACGACTACTATAGGGCGCCGGGAATTCGTCGACTG
GATCCGGTACCGAGGAGATCTGCCGCCGCGATCGCC
ATGTGGACAGCCAGAGGCCCTTCAGAAGACAGGTGGAGCAGTGAGGATGAGGAGGCTGCAGGGCCA
TCACAGGCTCTCTCCCCTCTACTTTCTGATACGCGCAAATTGTTTCTGAAGGAGAACTAGATCAGTTG
GCTCAGATTCGGCCATTAATATTTCAATTTTCATGAGCAGACAGCCATCAAGGATTGTTTGAAATCCTT
GAGGAAAAACAGCAGCGTATGATATCATGCAGGAATTTATGGCTTTAGAACTTAAGAATCTGCCTGGT
GAGTTCAACTCTGGGAATCAACCAAGCAACAGAGAAAAAACAGATACCGAGATATCTTCCATATGAT
TCAACACGCGTTCCTCTTGGAAAAAGCAAGGACTACATCAATGCTAGTTATATTAGAATAGTCAATTGT
GGAGAAGAGTATTTTATATCGCTACTCAAGGACCACTGCTGAGCACCATAGATGACTTTTGCCAAATG
GTGTTGGAAAAATAATTCAAATGTTATTGCCATGATAACCAAGAGAGATAGAAGGTGGAATTCAAATGC
TACCATTACTGGCCATTTCTCTGAAGAAGCCATTGGAATTGAAACACTCCCGTGTATTCTGGAGAAC
TACCAGATACTTCAATATTTTCATCATTGCAATGTTTCAAGTTGTGGAGAAGTCCACGGAACTAGTCAC
TCTGTAAACAGTTGCAGTTCACCAAGTGGCCAGACCATGGCACTCCTGCCTCAGCAGATAGCTTCATA
AAATATATTCGTTATGCAAGGAAGAGCCACCTTACAGGACCCATGGTTGTTCACTGCAGTGCCGGCATA
GGCCGGACAGGGGTGTTCTATGTGTGGATGTCGTGTTCTGTGCCATCGTAAAGAAGTTCATTCAAC
ATCATGGATATAGTGGCCAAATGAGAGAACAACGTTCTGGCATGTTCAAACGAAGGAGCAGTATCAC
TTTTGTTACGATATTGTGCTTGAAGTCTTCGGAACTTCTGACTTTGGATTAA
AGCGGACCGACGCGTACGCGGCCCTCGAGCAGAACTCATCTCAGAAGAGGATCTGGCAGCAAATGAT
ATCCTGGATTACAAGGATGACGACGATAAGGTTTAA
```

Restriction Sites: SgfI-RsrII



[View online »](#)

Plasmid Map:



ACCN: NM_001042395

Insert Size: 1020 bp

OTI Disclaimer: Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).

OTI Annotation: This TrueClone is provided through our Custom Cloning Process that includes sub-cloning into OriGene's pCMV6 vector and full sequencing to provide a non-variant match to the expected reference without frameshifts, and is delivered as lyophilized plasmid DNA.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method:

1. Centrifuge at 5,000xg for 5min.
2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
3. Close the tube and incubate for 10 minutes at room temperature.
4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

RefSeq: [NM_001042395.1](#)

RefSeq Size: 2872 bp

RefSeq ORF: 1020 bp

Locus ID: 653129

Cytogenetics: 10q11.22

Protein Families: Druggable Genome

MW: 39.2 kDa

Gene Summary: The product of this gene belongs to the family of classical tyrosine-specific protein tyrosine phosphatases. Many protein tyrosine phosphatases have been shown to regulate fundamental cellular processes and several are mutated in human diseases. Chromosome 10q contains a segmental duplication resulting in multiple copies of the protein tyrosine phosphatase, non-receptor type 20 gene. The two nearly identical copies are designated as PTPN20A and PTPN20B. A third copy is only partially duplicated and contains a pseudogene, designated as PTPN20C. This gene encodes the more centromeric copy, PTPN20A. Multiple alternatively spliced transcript variants encoding different isoforms have been identified. [provided by RefSeq, Jul 2008]

Transcript Variant: This variant (8) differs in the 5' UTR and in the 5' coding region, compared to variant 1. The resulting protein (isoform 8) contains a distinct N-terminus, compared to isoform 1.