

Product datasheet for **SC312368**

TBC1D20 (AK055488) Human Untagged Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	TBC1D20 (AK055488) Human Untagged Clone
Tag:	Tag Free
Symbol:	TBC1D20
Vector:	<u>pCMV6 series</u>
Fully Sequenced ORF:	>NCBI ORF sequence for AK055488, the custom clone sequence may differ by one or more nucleotides
Restriction Sites:	Please inquire
ACCN:	AK055488
OTI Disclaimer:	Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).
OTI Annotation:	This TrueClone is provided through our Custom Cloning Process that includes sub-cloning into OriGene's pCMV6 vector and full sequencing to provide a non-variant match to the expected reference without frameshifts, and is delivered as lyophilized plasmid DNA.
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	<ol style="list-style-type: none">1. Centrifuge at 5,000xg for 5min.2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.3. Close the tube and incubate for 10 minutes at room temperature.4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
RefSeq:	<u>AK055488.1</u>
RefSeq Size:	3218 bp
RefSeq ORF:	3218 bp

[View online »](#)

Locus ID: 128637

Cytogenetics: 20p13

Protein Families: Transmembrane

Gene Summary: This gene encodes a protein that belongs to a family of GTPase activator proteins of Rab-like small GTPases. The encoded protein and its cognate GTPase, Rab1, bind the nonstructural protein 5A (NS5A) of the hepatitis C virus (HCV) to mediate viral replication. Depletion of this protein inhibits replication of the virus and HCV infection. Mutations in this gene are associated with Warburg micro syndrome 4. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Mar 2014]