

Product datasheet for SC312466

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

CD89 (FCAR) (NM_133274) Human Untagged Clone

Product data:

Product Type: Expression Plasmids

Product Name: CD89 (FCAR) (NM_133274) Human Untagged Clone

Tag: Tag Free
Symbol: FCAR

Synonyms: CD89; CTB-61M7.2; FcalphaRI

Vector: pCMV6 series

Fully Sequenced ORF: >NCBI ORF sequence for NM_133274, the custom clone sequence may differ by one or more

nucleotides

CCTCGTGGCTCTCTTGGCCATACTGGT

Restriction Sites: Please inquire **ACCN:** NM 133274

OTI Disclaimer: Our molecular clone sequence data has been matched to the reference identifier above as a

point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative

RNA splicing form or single nucleotide polymorphism (SNP).

OTI Annotation: This TrueClone is provided through our Custom Cloning Process that includes sub-cloning

into OriGene's pCMV6 vector and full sequencing to provide a non-variant match to the expected reference without frameshifts, and is delivered as lyophilized plasmid DNA.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).





Reconstitution Method:

- 1. Centrifuge at 5,000xg for 5min.
- 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
- 3. Close the tube and incubate for 10 minutes at room temperature.
- 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
- 5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

RefSeq: <u>NM 133274.1</u>, <u>NP 579808.1</u>

 RefSeq Size:
 1561 bp

 RefSeq ORF:
 630 bp

 Locus ID:
 2204

 UniProt ID:
 P24071

 Cytogenetics:
 19q13.42

Protein Families: Transmembrane

Gene Summary: This gene is a member of the immunoglobulin gene superfamily and encodes a receptor for

the Fc region of IgA. The receptor is a transmembrane glycoprotein present on the surface of myeloid lineage cells such as neutrophils, monocytes, macrophages, and eosinophils, where it mediates immunologic responses to pathogens. It interacts with IgA-opsonized targets and triggers several immunologic defense processes, including phagocytosis, antibody-dependent cell-mediated cytotoxicity, and stimulation of the release of inflammatory mediators. Multiple alternatively spliced transcript variants encoding different isoforms have been described for

this gene. [provided by RefSeq, Jul 2008]

Transcript Variant: This variant (6, also known as U09) lacks an in-frame exon in the 5' coding region and uses an alternate splice site in the 3' coding region, which results in a frameshift, compared to variant 1. The encoded isoform (f) has a distinct C-terminus and is shorter than isoform a. Sequence Note: This RefSeq record was created from transcript and genomic sequence data to make the sequence consistent with the reference genome assembly. The genomic coordinates used for the transcript record were based on transcript alignments.