

## **Product datasheet for SC312551**

## SS18 (AK098354) Human Untagged Clone

## **Product data:**

**Product Type:** Expression Plasmids

**Product Name:** SS18 (AK098354) Human Untagged Clone

Tag: Tag Free Symbol: SS18

Synonyms: PACS-2; PACS1L Vector: pCMV6 series

Fully Sequenced ORF: >NCBI ORF sequence for AK098354, the custom clone sequence may differ by one or more

nucleotides

GGGGGAGAGGTAGGAAGGTCTCCCAT

**Restriction Sites:** Please inquire ACCN: AK098354

**OTI Disclaimer:** Our molecular clone sequence data has been matched to the reference identifier above as a

point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative

RNA splicing form or single nucleotide polymorphism (SNP).

OTI Annotation: This TrueClone is provided through our Custom Cloning Process that includes sub-cloning

into OriGene's pCMV6 vector and full sequencing to provide a non-variant match to the expected reference without frameshifts, and is delivered as lyophilized plasmid DNA.

**Components:** The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).



**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



**Reconstitution Method:** 

- 1. Centrifuge at 5,000xg for 5min.
- 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
- 3. Close the tube and incubate for 10 minutes at room temperature.
- 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
- 5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

**RefSeq:** <u>AK098354.1</u>, <u>BAC05290.1</u>

RefSeq Size: 2663 bp
RefSeq ORF: 690 bp
Locus ID: 6760
Cytogenetics: 18q11.2

**Protein Families:** Druggable Genome

**Gene Summary:** Appears to function synergistically with RBM14 as a transcriptional coactivator. Isoform 1 and

isoform 2 function in nuclear receptor coactivation. Isoform 1 and isoform 2 function in general transcriptional coactivation. Component of SWI/SNF chromatin remodeling

subcomplex GBAF that carries out key enzymatic activities, changing chromatin structure by

altering DNA-histone contacts within a nucleosome in an ATP-dependent manner

(PubMed:29374058).[UniProtKB/Swiss-Prot Function]