

Product datasheet for **SC313004**

C9orf140 (SAPCD2) (BC009435) Human Untagged Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	C9orf140 (SAPCD2) (BC009435) Human Untagged Clone
Tag:	Tag Free
Symbol:	C9orf140
Synonyms:	2010317E24Rik; chromosome 9 open reading frame 140; OTTHUMP00000022651; p42.3; tumor specificity and mitosis phase-dependent expression protein
Vector:	<u>pCMV6 series</u>
Fully Sequenced ORF:	>NCBI ORF sequence for BC009435, the custom clone sequence may differ by one or more nucleotides
Restriction Sites:	Please inquire
ACCN:	BC009435
OTI Disclaimer:	Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).
OTI Annotation:	This TrueClone is provided through our Custom Cloning Process that includes sub-cloning into OriGene's pCMV6 vector and full sequencing to provide a non-variant match to the expected reference without frameshifts, and is delivered as lyophilized plasmid DNA.
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	<ol style="list-style-type: none"> 1. Centrifuge at 5,000xg for 5min. 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA. 3. Close the tube and incubate for 10 minutes at room temperature. 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom. 5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
RefSeq:	<u>BC009435.1</u> , <u>AAH09435.1</u>
RefSeq Size:	2472 bp


[View online »](#)

RefSeq ORF: 2472 bp

Locus ID: 89958

Cytogenetics: 9q34.3

Gene Summary: Plays a role in planar mitotic spindle orientation in retinal progenitor cells (RPCs) and promotes the production of symmetric terminal divisions (By similarity). Negatively regulates the mitotic apical cortex localization of GPSM2 (PubMed:26766442). Involved also in positive regulation of cell proliferation and tumor cell growth (PubMed:23576022, PubMed:23704824). [UniProtKB/Swiss-Prot Function]