

## Product datasheet for SC313098

### ZDHHC20 (NM\_153251) Human Untagged Clone

#### Product data:


Product Type:	Expression Plasmids
Product Name:	ZDHHC20 (NM_153251) Human Untagged Clone
Tag:	Tag Free
Symbol:	ZDHHC20
Synonyms:	4933421L13Rik; DHHC-20; DHHC20
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)
Fully Sequenced ORF:	>SC313098 representing NM_153251. Blue=Insert sequence Red=Cloning site Green=Tag(s)

```
GCTCGTTTGTAGTGAACCGTCAGAATTTTGTAAACGACTCACTATAGGGCGCCGGGAATTCGTCGACTG
GATCCGGTACCGAGGAGATCTGCCGCCGCGATCGCC
ATGGCGCCTGGACGCTGTGGCGCTGCTGCCAGCGCGTCGTGGGCTGGGTGCCGGTGCTCTTCATCACC
TTCGTGGTCTGCTGGTCTACTACGCGTACGTGGTGGAGCTCTGCGTGTTACTATTTTTGGAAATGAA
GAAAATGGAAAGACCGTTGTTTACCTTGTGGCTTTCCATCTGTTCTTTGTTATGTTTGTATGGTCTAT
TGGATGACAATTTTACATCTCCCGCTTCCCGTCCAAAGAGTTCTACTGTCCAATTCTGAAAAGGAA
CGTTATGAAAAAAGAAATTCAGCCAAGAAAGACAACAAGAAATTTGAGAAGAGCAGCAAGAGCTTTACCT
ATCTATACCACATCAGCTTCAAAAATATCAGATATTGTGAAAAATGTCAGCTGATTAACCTGATCGG
GCGCATCACTGCTCAGCCTGTGACTCATGTATTCTTAAGATGGATCATCACTGTCCTTGGGTGAATAAC
TGTGTGGGATTTTCTAATTACAATTTCTTCTGCTGTTTTTATTGTATTCCCTATTATATTGCTTTTC
GTGGCTGCAACAGTTTTAGAGTACTTTATAAAAATTTGGACGAATGAACTGACAGATACAGTGCAGAAA
TTCCACGTAATTTTTCTTTCTTTGTGTCTGCAATGTTCTTTCATCAGCGTCTCTCACTTTTCAGCTAC
CACTGCTGGCTAGTTGGAAAAATAGAACAAATAGAATCATTCCGCGCACCCACGTTTTTCATACGGA
CCTGATGGAAATGGTTTCTCTCTTGGATGCAGTAAAAATGGAGACAAGTCTTTGGTGATGAAAAGAAA
TATTGGCTACTTCCAATATTTCAAGCTTGGGTGATGGTTGCAGTTTTCCAACCTGCCTTGTGGGGATG
GATCCAGAACAAAGCTTCTGTTACAAACCAGAATGAGTATGCCAGAAAGTGGCTCAAATCAACCTTTTCT
ATCAAACCACTTAGTGAATCAAAAACCGCTTGTGGACAGTGAATCTCAGTGGCTGGAGAAATGGAGCT
GAAGAAGGCATCGTCAAATCAGGTGATGA
AGCGGACCGACGCGTACGCGGCCGCTCGAGCAGAACTCATCTCAGAAGAGGATCTGGCAGCAAATGAT
ATCCTGGATTACAAGGATGACGACGATAAGGTTTAA
```

Restriction Sites: SgfI-RsrII



[View online »](#)

**Plasmid Map:**


**ACCN:** NM\_153251

**Insert Size:** 1065 bp

**OTI Disclaimer:** Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).

**OTI Annotation:** This TrueClone is provided through our Custom Cloning Process that includes sub-cloning into OriGene's pCMV6 vector and full sequencing to provide a non-variant match to the expected reference without frameshifts, and is delivered as lyophilized plasmid DNA.

**Components:** The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

**Reconstitution Method:**

1. Centrifuge at 5,000xg for 5min.
2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
3. Close the tube and incubate for 10 minutes at room temperature.
4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

**RefSeq:** [NM\\_153251.3](#)

**RefSeq Size:** 5356 bp

**RefSeq ORF:** 1065 bp

**Locus ID:** 253832

UniProt ID:	<u>Q5W0Z9</u>
Cytogenetics:	13q12.11
Protein Families:	Transmembrane
MW:	41.1 kDa
Gene Summary:	<p>Catalyzes palmitoylation of Cys residues on target proteins (PubMed:27153536, PubMed:29326245). Catalyzes palmitoylation of Cys residues in the cytoplasmic C-terminus of EGFR, and modulates the duration of EGFR signaling by modulating palmitoylation-dependent EGFR internalization and degradation (PubMed:27153536). Has a preference for acyl-CoA with C16 fatty acid chains (PubMed:29326245). Can also utilize acyl-CoA with C14 and C18 fatty acid chains (PubMed:29326245).[UniProtKB/Swiss-Prot Function]</p> <p>Transcript Variant: This variant (1) encodes the longer isoform (1). Sequence Note: This RefSeq record was created from transcript and genomic sequence data to make the sequence consistent with the reference genome assembly. The genomic coordinates used for the transcript record were based on transcript alignments.</p>