

Product datasheet for SC313294

Cytohesin 1 (CYTH1) (NM_017456) Human Untagged Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	Cytohesin 1 (CYTH1) (NM_017456) Human Untagged Clone
Tag:	Tag Free
Symbol:	Cytohesin 1
Synonyms:	B2-1; CYTOHESIN-1; D17S811E; PSCD1; SEC7
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)
Fully Sequenced ORF:	>SC313294 representing NM_017456. Blue=Insert sequence Red=Cloning site Green=Tag(s)

GCTCGTTTGTAGTAACCGTCAGAATTTTGTAAACGACTCACTATAGGGCGGCCGGGAATTCGTCGACTG
 GATCCGGTACCGAGGAGATCTGCCGCC**CGATCGCC**
 ATGGAGGAGGACGACAGCTACGTTCCAGTGACCTGACAGCAGAGGAGCGTCAAGAACTGGAGAACATC
 CGACGGAGAAAACAGGAGCTGCTGGCTGACATTCAGAGGCTGAAGGATGAGATAGCAGAAGTAGCTAAT
 GAAATTGAAAACCTGGGATCCACAGAGGAAAGGAAAAACATGCAGAGGAACAAACAGGTAGCCATGGGC
 AGGAAAAAATTTAATATGGACCTAAAAAGGGGATCCAGTTCTTAATAGAGAACGACCTCCTGAAGAAC
 ACTTGTGAAGACATTGCCAGTTCTTATATAAAGGCGAAGGGCTCAACAAGACAGCCATCGGCGACTAC
 CTAGGGGAGAGAGATGAGTTTAATATCCAGTTCTTCATGCATTTGTGGAGCTGCATGAGTTCAGTGAT
 CTTAATCTCGTCCAGGCACTACGGCAGTTCTGTGGAGCTTCCGGCTACCCGGAGAGGCCAGAAGATC
 GACCGGATGATGGAGGCGTTTGCCAGCGATATTGTCAGTGCAATAATGGCGTGTTCAGTCCACGGAT
 ACTTGTACGTCCTCTCCTTTGCCATCATCATGTTGAACACCAAGTCTGCACAACCCCAATGTCAAAGAT
 AAGCCCACTGTGGAGAGGTTCAATTGCCATGAACCGAGGCATCAATGATGGGGGAGACCTGCCGGAGGAG
 CTCCTCCGAATCTCTATGAGAGCATAAAAAATGAACCTTTAAATCCCAGAAGACGACGGGAATGAC
 CTCACTCACACTTTCTCAATCCAGACCGAGAAGGCTGGCTATTGAAACTCGGTGGCAGGGTAAAGACT
 TGAAGAGACGCTGGTTCACTGACTGACAACCTGCCTTTACTACTTTGAGTATACCACGGATAAGGAG
 CCCCCTGGAATCATCCCTTTAGAGAATCTGAGTATCCGGGAAGTGGAGGACTCCAAAAACCAAATGCTG
 TTTGAGCTTTATATCCCGACAATAAAGACCAAGTTATCAAGGCCTGCAAGACCGAGGCTGACGGGCGG
 GTGGTGGAGGGGAACCACTGTTTACCGGATCTCAGCTCCGACGCCCGAGGAGAAGGAGGAGTGGATT
 AAGTGCATTAAGCAGCCATCAGCAGGGACCTTTCTACGAAATGCTCGCAGCACGGAAGAAAGAGGTC
 TCCTCCACGAAGCGACACTGA
 ACGCGTACGCGGCCGCTCGAGCAGAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGAT
 TACAAGGATGACGACGATAAGGTTTAAACGGCCGCGC

Restriction Sites: SgfI-MluI


[View online »](#)

ACCN:	NM_017456
Insert Size:	1194 bp
OTI Disclaimer:	Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).
OTI Annotation:	This TrueClone is provided through our Custom Cloning Process that includes sub-cloning into OriGene's pCMV6 vector and full sequencing to provide a non-variant match to the expected reference without frameshifts, and is delivered as lyophilized plasmid DNA.
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	<ol style="list-style-type: none"> 1. Centrifuge at 5,000xg for 5min. 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA. 3. Close the tube and incubate for 10 minutes at room temperature. 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom. 5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
RefSeq:	<u>NM_017456.3</u>
RefSeq Size:	3363 bp
RefSeq ORF:	1194 bp
Locus ID:	9267
UniProt ID:	<u>Q15438</u>
Cytogenetics:	17q25.3
Domains:	Sec7, PH
MW:	46.4 kDa

Gene Summary:

The protein encoded by this gene is a member of the PSCD family. Members of this family have identical structural organization that consists of an N-terminal coiled-coil motif, a central Sec7 domain, and a C-terminal pleckstrin homology (PH) domain. The coiled-coil motif is involved in homodimerization, the Sec7 domain contains guanine-nucleotide exchange protein activity, and the PH domain interacts with phospholipids and is responsible for association of PSCDs with membranes. Members of this family appear to mediate the regulation of protein sorting and membrane trafficking. This gene is highly expressed in natural killer and peripheral T cells, and regulates the adhesiveness of integrins at the plasma membrane of lymphocytes. A pseudogene of this gene has been defined on the X chromosome. Alternative splicing results in multiple transcript variants. [provided by RefSeq, May 2014]

Transcript Variant: This variant (2) lacks an in-frame exon in the central coding region, compared to variant 1. The encoded isoform (2) is shorter, compared to isoform 1.