

Product datasheet for SC313551

NARF (NM_012336) Human Untagged Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	NARF (NM_012336) Human Untagged Clone
Tag:	Tag Free
Symbol:	NARF
Synonyms:	IOP2
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)
Fully Sequenced ORF:	>SC313551 representing NM_012336. Blue=Insert sequence Red=Cloning site Green=Tag(s)

```

GCTCGTTTGTAGTGAACCGTCAGAATTTTGTAAATACGACTCACTATAGGGCGGCCGGGAATTCGTCGACTG
GATCCGGTACCGAGGAGATCTGCCGCCGCGATCGCC
ATGAAGTGTGAGCACTGCACGCGCAAGGAATGTAGTAAGAAAACAAAAGTATGACCAAGAGAATGTG
TCAGCCGATGCACCGAGTCCAGCCCAGGAAAATGGAGAGAAGGGAGAATTCACAAAGTTGGCTGATGCC
AAGATATTTTGTAGCGACTGCCTGGCATGTGACAGCTGTATGACTGCAGAGGAAGGAGTCCAACCTTTCC
CAGCAAAATGCCAAGGACTTCTTCGCGTTCTGAACCTTAACAAGAAATGTGATACCTCAAAGCACAAA
GTGCTGGTAGTGTCTGTGTCTCAATCTTTGCCTTATTTTGTCTAAATTCAACCTCAGTGTAAC
GATGCATCCAGAAGACTCTGTGGTTTCTCAAAAGTCTTGGGGTGCACTATGTATTTGATACGACGATA
GCTGCGGATTTTAGTATCCTGGAGAGTCAAAAAGAATTCGTGCGTCGCTATCGCCAGCACAGTGAGGAG
GAACGCACCCTGCCCATGCTGACCTCTGCTGTCTGGTGGGTCCGATACGCCGAGCGGGTGCTGGGT
CGCCCCATCACTGCCACCTCTGCACCGCAAGTCCCCCAGCAGGTGATGGGCTCTTTGGTGAAGGAT
TATTTGCGCAGACAGCAGAACCTGTCTCCAGAGAAGATTTCCACGTGATTGTGGCCCTTGTATGAC
AAGAAGCTGGAGGCTCTTCAGGAAAGCCTTCCCCTGCTTTGCATGGCTCCCGGGGCGCTGACTGCGTG
TTAACATCAGGTGAAATTGCTCAAATAATGGAGCAAGGTGACCTCTCAGTGAGAGATGCTGCCGTCGAC
ACTCTGTTTGGAGACTTGAAGGAGGACAAAGTGACGCGTCATGATGGAGCCAGCTCAGACGGGCACCTG
GCACACATCTTCAGACATGCGGCCAAGGAGCTGTTCAACGAGGATGTGGAGGAGTCACTTACCGAGCC
CTGAGAAACAAAGACTTCCAAGAGGTCAACCTTGAGAAGAACGGAGAGGTGGTGTACGCTTTGCTGCA
GCCTATGGCTTTGCAAAACATCCAGAACATGATCCTGAAGCTTAAGAAGGGCAAGTTCCTATTCCACTTT
GTGGAGGTCTCGCTGTGCTGGAGGATGCTTAAATGGCAGAGGCCAAGCCAGACTCCAGACGGACAT
GCGGATAAGGCCCTGCTGCGGCAGATGGAAGGCATTTACGCTGACATCCCTGTGCGGCGTCCGGAGTCC
AGTGACACAGTGCAGGAGCTGTACCAGGAGTGGCTGGAGGGGATCAACTCCCCAAGGCCGAGAGGTG
CTGCATACCAGTACCAGAGCCAGGAGCGTGGCACACACAGCCTGGACATCAAGTGGTGA
AGCGGACCGACGCTACGCGGCCGCTCGAGCAGAACTCATCTCAGAAGAGGATCTGGCAGCAAATGAT
ATCCTGGATTACAAGGATGACGACGATAAGGTTTAA

```


[View online »](#)

Restriction Sites:	Sgfl-RsrII
ACCN:	NM_012336
Insert Size:	1371 bp
OTI Disclaimer:	Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).
OTI Annotation:	This TrueClone is provided through our Custom Cloning Process that includes sub-cloning into OriGene's pCMV6 vector and full sequencing to provide a non-variant match to the expected reference without frameshifts, and is delivered as lyophilized plasmid DNA.
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	<ol style="list-style-type: none"> 1. Centrifuge at 5,000xg for 5min. 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA. 3. Close the tube and incubate for 10 minutes at room temperature. 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom. 5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
RefSeq:	<u>NM_012336.3</u>
RefSeq Size:	1638 bp
RefSeq ORF:	1371 bp
Locus ID:	26502
UniProt ID:	<u>Q9UHQ1</u>
Cytogenetics:	17q25.3
Domains:	Fe_hyd_SSU, Fe_hyd_Ig_C
MW:	51.2 kDa

Gene Summary:

Several proteins have been found to be prenylated and methylated at their carboxyl-terminal ends. Prenylation was initially believed to be important only for membrane attachment. However, another role for prenylation appears to be its importance in protein-protein interactions. The only nuclear proteins known to be prenylated in mammalian cells are prelamin A- and B-type lamins. Prelamin A is farnesylated and carboxymethylated on the cysteine residue of a carboxyl-terminal CaaX motif. This post-translationally modified cysteine residue is removed from prelamin A when it is endoproteolytically processed into mature lamin A. The protein encoded by this gene binds to the prenylated prelamin A carboxyl-terminal tail domain. It may be a component of a prelamin A endoprotease complex. The encoded protein is located in the nucleus, where it partially colocalizes with the nuclear lamina. It shares limited sequence similarity with iron-only bacterial hydrogenases. Alternatively spliced transcript variants encoding different isoforms have been identified for this gene, including one with a novel exon that is generated by RNA editing. [provided by RefSeq, Jul 2008]

Transcript Variant: This variant (1) encodes the predominant isoform (a).