

## Product datasheet for **SC314670**

### Cullin 1 (CUL1) (AF062536) Human Untagged Clone

#### Product data:

Product Type:	Expression Plasmids
Product Name:	Cullin 1 (CUL1) (AF062536) Human Untagged Clone
Tag:	Tag Free
Symbol:	Cullin 1
Vector:	<u><a href="#">pCMV6 series</a></u>



[View online »](#)

**Fully Sequenced ORF:** >NCBI ORF sequence for AF062536, the custom clone sequence may differ by one or more nucleotides

```

ATGTCGTC AACCCGGAGCCAGAACCCCCACGGCCTGAAGCAGATTGGCCTGGACCAGATC
TGGGACGACCTCAGAGCCGGCATCCAGCAGGTGTACACACGGCAGAGCATGGCCAAGTCC
AGATATATGGAGCTCTACACTCATGTTTATAACTACTGTACTAGTGTCCACCAGTCAAAC
CAAGCACGAGGAGCTGGAGTTCCTCCTTCTAAGTCGAAAAAGGGGAGACACCTGGAGGA
GCTCAGTTTGTGGCCTGGAATTATATAAACGACTTAAGGAATTTTTGAAGAATTACTTG
ACAAATCTTCTTAAGGATGGAGAAGATTTGATGGATGAGAGTGTACTGAAATTCTACACT
CAACAATGGGAAGATTATCGATTTTCAAGCAAAGTGCTGAATGGAATTTGTGCCTACCTC
AATAGACATTGGGTTTCGCCGTGAATGTGACGAAGGACGAAAAGGAATATATGAAATCTAT
TCGCTTGCAATTGGTGACTTGGAGAGACTGTCTGTTGAGGCCACTGAATAAACAGGTAACA
AATGCTGTTTTAAAGCTGATTGAAAAGGAAAGGAATGGTGAACCATCAATACAAGATTG
ATTAGTGGAGTTGTACAGTCTTACGTGGAATGGGGCTGAATGAAGATGATGCATTTGCA
AAGGGCCCTACGTTAACAGTGTATAAAGAATCCTTTGAATCTCAATTTTTGGCTGACACA
GAGAGATTTTATACCAGAGAGAGTACTGAATTCTGACGACGAAACCCAGTTACTGAATAT
ATGAAAAAGGCAGAGGCTCGTCTGCTTGAGGAACAACGAAGAGTTCAGGTTTACCTTCAT
GAAAGCACACAAGATGAATTAGCAAGGAAATGTGAACAAGTCTCATTGAAAAACTTTG
GAAATTTTCCACACAGAAATTTGAGAATTTATTGGATGCTGACAAAAATGAAGATTTGGGA
CGCATGTATAATCTTGTATCTAGAATCCAGGATGGCCTAGGAGAAATGAAAAACTGTTG
GAGACACACATTCATAATCAGGGTCTTGACGCCATTGAAAAGTGTGGAGAAGCTGCTTTA
AATGACCCCAAAATGTATGTACAGACAGTGTCTGATGTTCAAAAAATACAATGCCCTG
GTAATGTCTGCATTCAACAATGACGCTGGCTTTGTGGCTGCTCTTGATAAGGCTTGTGGT
CGTTCATAAACAACAACGCGGTTACCAAGATGGCCCAATCATCCAGTAAATCCCCTGAG
TTGCTGGCTCGATACTGTGACTCCTTGTGAAGAAAAGTTCCAAGAACCAGAGGAGGCA
GAACTAGAAGACACACTCAATCAAGTGATGGTTGTCTTCAAGTACATAGAAGACAAGAC
GTATTTGAGAAGTTCTATGCGAAGATGCTCGCCAAGAGGCTCGTCCACCAGAACAGTGCA
AGTGACGATGCCGAAGCCAGCATGATCTCCAAGTTAAAGCAAGCTTGC GGTTTCGAGTAC
ACCTCTAAACTTCAGCGCATGTTTCAAGACATTGGCGTGAGCAAAGATCTGAACGAGCAA
TTCAAAAAGCACTTGACAAACTCAGAACCCTAGACTTGGATTTTCAAGTCAAGTGCTG
AGCTCCGGGCTCGGCCCTTCCAGCAGTCTGTACATTTGCCTTGCCGTGAGAGTTGGAA
CGTAGTTATCAGCGATTCACAGTCTTCTACGCCAGCCGCACAGTGGCCGAAAATTGACG
TGTTTATATCAGTTGTCTAAAGGAGAATTGGTAACTAAGTCTTCAAAAACAGATATACT
TTGACGGCGTCGACATTCAGATGGCTATCCTGCTTCAAGTACAACACGGAAGATGCCTAC
ACTGTGCAGCAGCTGACCAGCAGCACTCAATTAATAATGGACATTTTGGCGCAAGTTTTA
CAGATTTTATTAAGTCGAAGCTATTGGTCTTGGAAAGTGAATGCAAAATGTTGATGAG
GTGGAATTGAAGCCAGATACCTTAATAAAATTATATCTTGGTTATAAAAAAAGAATAA
AGGGTTAACATCAATGTGCCAATGAAAACCGAACAGAAGCAGGAACAAGAAACACACAC
AAAAACATCGAGGAAGACCGCAAACACTGATTTCAGGCGGCCATCGTGAATCATGAAG
ATGAGGAAGGTTCTGAAACACCAGCAGTTACTTGGCGAGGTCCTCACTCAGCTGTCTCC
AGGTTCAAACCTCGAGTCCCTGTGATCAAGAAATGCATTGACATTCTAATTGAGAAAGAA
TATTTGGAGCGAGTGGATGGTAAAAGGACACCTACAGTTACTTGGCT

```

**Restriction Sites:** Please inquire

**ACCN:** AF062536

**OTI Disclaimer:** Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).

<b>OTI Annotation:</b>	This TrueClone is provided through our Custom Cloning Process that includes sub-cloning into OriGene's pCMV6 vector and full sequencing to provide a non-variant match to the expected reference without frameshifts, and is delivered as lyophilized plasmid DNA.
<b>Components:</b>	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
<b>Reconstitution Method:</b>	<ol style="list-style-type: none"> <li>1. Centrifuge at 5,000xg for 5min.</li> <li>2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.</li> <li>3. Close the tube and incubate for 10 minutes at room temperature.</li> <li>4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.</li> <li>5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.</li> </ol>
<b>RefSeq:</b>	<a href="#">AF062536.1</a> , <a href="#">AAC36681.1</a>
<b>RefSeq Size:</b>	2583 bp
<b>RefSeq ORF:</b>	2583 bp
<b>Locus ID:</b>	8454
<b>Cytogenetics:</b>	7q36.1
<b>Domains:</b>	CULLIN
<b>Protein Families:</b>	Druggable Genome
<b>Protein Pathways:</b>	Cell cycle, Oocyte meiosis, TGF-beta signaling pathway, Ubiquitin mediated proteolysis, Wnt signaling pathway

**Gene Summary:**

Core component of multiple cullin-RING-based SCF (SKP1-CUL1-F-box protein) E3 ubiquitin-protein ligase complexes, which mediate the ubiquitination of proteins involved in cell cycle progression, signal transduction and transcription. SCF complexes and ARIH1 collaborate in tandem to mediate ubiquitination of target proteins (PubMed:27565346). In the SCF complex, serves as a rigid scaffold that organizes the SKP1-F-box protein and RBX1 subunits. May contribute to catalysis through positioning of the substrate and the ubiquitin-conjugating enzyme. The E3 ubiquitin-protein ligase activity of the complex is dependent on the neddylation of the cullin subunit and exchange of the substrate recognition component is mediated by TIP120A/CAND1. The functional specificity of the SCF complex depends on the F-box protein as substrate recognition component. SCF(BTRC) and SCF(FBXW11) direct ubiquitination of CTNNB1 and participate in Wnt signaling. SCF(FBXW11) directs ubiquitination of phosphorylated NFKBIA. SCF(BTRC) directs ubiquitination of NFKBIB, NFKBIE, ATF4, SMAD3, SMAD4, CDC25A, FBXO5 and probably NFKB2. SCF(BTRC) and/or SCF(FBXW11) direct ubiquitination of CEP68 (PubMed:25704143, PubMed:25503564). SCF(SKP2) directs ubiquitination of phosphorylated CDKN1B/p27kip and is involved in regulation of G1/S transition. SCF(SKP2) directs ubiquitination of ORC1, CDT1, RBL2, ELF4, CDKN1A, RAG2, FOXO1A, and probably MYC and TAL1. SCF(FBXW7) directs ubiquitination of CCNE1, NOTCH1 released notch intracellular domain (NICD), and probably PSEN1. SCF(FBXW2) directs ubiquitination of GCM1. SCF(FBXO32) directs ubiquitination of MYOD1. SCF(FBXO7) directs ubiquitination of BIRC2 and DLGAP5. SCF(FBXO33) directs ubiquitination of YBX1. SCF(FBXO1) directs ubiquitination of BCL6 and DTL but does not seem to direct ubiquitination of TP53. SCF(BTRC) mediates the ubiquitination of NFKBIA at 'Lys-21' and 'Lys-22'; the degradation frees the associated NFKB1-RELA dimer to translocate into the nucleus and to activate transcription. SCF(CCNF) directs ubiquitination of CCP110. SCF(FBXL3) and SCF(FBXL21) direct ubiquitination of CRY1 and CRY2. SCF(FBXO9) directs ubiquitination of TT11 and TELO2. SCF(FBXO10) directs ubiquitination of BCL2.[UniProtKB/Swiss-Prot Function]