

Product datasheet for SC314822

PARP9 (NM_031458) Human Untagged Clone

Product data:

Product Type: Expression Plasmids
Product Name: PARP9 (NM_031458) Human Untagged Clone
Tag: Tag Free
Symbol: PARP9
Synonyms: ARTD9; BAL; BAL1; MGC:7868
Mammalian Cell Selection: None
Vector: pCMV6-XL4
E. coli Selection: Ampicillin (100 ug/mL)

Fully Sequenced ORF: >OriGene sequence for NM_031458 edited
CGGCCGCGCGCCTGCCGGAAGTGGGCCACCATATCTGGAACTACAGTCTATGCTTTGA
AGCGAAAAGGGAATAAACATTTAAAGACTCCCCGGGGACCTGGAGGATGGACTTTTCC
ATGGTGGCCGGAGCAGCAGCTTACAATGAAAAATCAGGTAGGATTACCTCGCTCTCACTC
TTGTTTCAGAAAGTCTTTGCTCAGATCTTCTCAGTGGAGAAAGGGGAATACAGAAGAA
TGTCTCCCTACAAGTGCTCAGAGACTGGTCTTGGAGAAACTATAGTTGGCAAATT
CCCATTAAACCACAATGACTTCAAATTTTAAAAAATAATGAGCGTCAGCTGTGTGAAGTC
CTCCAGAATAAGTTTGGCTGTATCTCTACCCTGGTCTCTCCAGTTCAGGAAGGCAACAGC
AAATCTCTGCAAGTGTTCAGAAAAATGCTGACTCTAGGATAGAGTTATCAGTCTGGAAA
GATGACCTCACCACACATGCTGTTGATGCTGTGGTGAATGCAGCCAATGAAGATCTTCTG
CATGGGGGAGGCCTGGCCCTGGCCTGGTAAAAGCTGGTGGATTTGAAATCCAAGAAGAG
AGCAAACAGTTTGTGCCAGATATGGTAAAGTGTGAGCTGGTGAATAGCTGTCACGGGA
GCAGGGAGGCTTCCCTGCAAACAGATCATCCATGCTGTTGGGCCTCGGTGGATGGAATGG
GATAAACAGGGATGTAAGTGGAAAGCTGCAGAGGGCCATTGTAAGTATTCTGAATTATGTC
ATCTATAAAAAACTCACATTAAGACAGTAGCAATTCAGCCTTGAGCTCTGGGATTTTT
CAGTTCCTCTGAATTTGTGACAAAGACTATTGTAGAGACTATCCGGTTAGTTTGCAA
GGGAAGCCAATGATGAGTAATTTGAAAGAAATTCACCTGGTGAAGTGAAGTGGGACAAGAA
ACCACCCCTTTTCAATGCAATGGTGTGAAACAACCTGACCCCTCCAGATTGTCAGGGC
CACATTTGAATGGCAGACGCAGATGTAATTGTTAATTCTGTAAACCCACATGATATTACA
GTTGGACCTGTGGCAAAGTCAATTTCTACAACAAGCAGGAGTTGAAATGAAATCGGAATTT
CTTGCCACAAAGGCTAAACAGTTTCAACGGTCCCAGTTGGTACTGGTCACAAAAGGATTT
AACTTGTCTGTAATATATATACCATGTAAGTGGCATTGAGAAATTTCTAAACCTCAG
ATATTAACATGCAATGAAGGAGTGTGGAAAAATGCATTGAGCAAAATATAACTTCC
ATTTCTTCTGCCCCTGGGACTGGAACATGGAATAAAGAAGGAAACAGCAGCAGAG
ATTTTGTGATGAAGTTTTAACATTTGCCAAAGACCATGTAACACACAGTTAACTGTA
AAATTTGTGATCTTCCAACAGATTTGGAGATATATAAGGCTTTCAGTTCTGAAATGGCA



[View online »](#)

```

AAGAGGTCCAAGATGCTGAGTTTGAACAATTACAGTGTCCCCAGTCAACCAGAGAGGAG
AAAAGAGAAAAATGGGCTTGAAGCTAGATCTCCTGCCATCAATCTGATGGGATTCAACGTG
GAAGAGATGTATGAGGCCACGCATGGATCCAAAGAATCCTGAGTCTCCAGAACCACCAC
ATCATTGAGAATAATCATATTCTGTACCTTGGGAGAAAGGAACATGACATTTTGTCTCAG
CTTCAGAAAACCTCAAGTGTCTCCATCACAGAAATATCAGCCCAGGAAGGACAGAGTTA
GAGATTGAAGGAGCCCGGGCTGACCTCATTGAGGTGGTTATGAACATTGAAGATATGCTT
TGTAAGTACAGGAGGAAATGGCAAGGAAAAAGGAGCGAGGCCTTTGGCGCTCGTTAGGA
CAGTGGACTATTAGCAACAAAAACCCAAGACGAAATGAAAGAAAATATCATATTTCTG
AAATGTCTGTGCCTCCAACCTCAAGAGCTTCTAGATCAAAAGAAAACAGTTTGAAAAATGT
GGTTTGCAGGTTCTAAAGGTGGAGAAGATAGACAATGAGGTCTTATGGCTGCCTTTCAA
AGAAAGAAGAAAATGATGGAAGAAAACTGCACAGGCAACCTGTGAGCCATAGGCTGTTT
CAGCAAGTCCCATACCAGTTCTGCAATGTGGTATGCAGAGTTGGCTTTCAAAGAATGTAC
TCGACACCTTGCATCCAAAATACGGAGCTGGCATATACTTCACCAAGAACCTCAAAAAC
CTGGCAGAGAAGCCAAGAAAATCTCTGCTGCAGATAAGCTGATCTATGTGTTTGAGGCT
GAAGTACTCACAGGCTTCTTCTGCCAGGACATCCGTTAAATATTGTTCCCCCACCCTG
AGTCTGGAGCTATAGATGGTACATGACAGTGTGGTTGACAATGTCTCCAGCCCTGAAACC
TTTGTTATTTTTAGTGGCATGCAGGCTATACCTCAGTATTTGTGGACATGCACCCAGGAA
TATGTACAGTCACAAGATTACTCATCAGGACCAATGAGACCCTTTGCACAGCATCCTTGG
AGGGGATTCGCAAGTGGCAGCCCTGTTGATTAATCTCTACATCATTTTAAACAGCTGGTAT
GGCCTTACCTTGGGTGAACAAATAATGACCATCGATGGCTCAAAGAGTGGCTTGA
ATATATCCCATGGGTTATCTGTATGGACTGACTGGGTTATTGAAAGGACTAGCCACATAC
TAGCATCTTAGTCCCTTATCTGTCTTTATGTCTTGGGGTTGGGTAGGTAGATACAAA
TGAAACACTTTTCAGGACCTTCTCTCTCTTGCAGTTGTTCTTTAATCTCCTTTACTAGAG
GAGATAAATATTTTGCATATAAATGAAGAAATTTTCTAGTATATAACGCAGGCCTTTTAT
TTTCTAAAATGATGATAGTATAAAAATGTTAGGATAACAGAATGATTTTATGATTTTCCAG
AGAATATTATAAAGTGCTTTAGGTATGAAAATAAATCATCTTTGTCTGATTAAAAAAA
AAA
    
```

- Restriction Sites:** NotI-NotI
- ACCN:** NM_031458
- Insert Size:** 3100 bp
- OTI Disclaimer:** Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).
- OTI Annotation:** The insert of this clone has been fully sequenced and found to be a perfect match to NM_031458.1 except for one SNP.
- Components:** The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
- Reconstitution Method:**
1. Centrifuge at 5,000xg for 5min.
 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
 3. Close the tube and incubate for 10 minutes at room temperature.
 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
 5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

RefSeq: [NM_031458.1](#), [NP_113646.1](#)

RefSeq Size: 3243 bp

RefSeq ORF: 2565 bp

Locus ID: 83666

UniProt ID: [Q8IXQ6](#)

Cytogenetics: 3q21.1

Domains: A1pp

Gene Summary: ADP-ribosyltransferase which, in association with E3 ligase DTX3L, plays a role in DNA damage repair and in immune responses including interferon-mediated antiviral defenses (PubMed:16809771, PubMed:23230272, PubMed:26479788, PubMed:27796300). Within the complex, enhances DTX3L E3 ligase activity which is further enhanced by PARP9 binding to poly(ADP-ribose) (PubMed:28525742). In association with DTX3L and in presence of E1 and E2 enzymes, mediates NAD(+)-dependent mono-ADP-ribosylation of ubiquitin which prevents ubiquitin conjugation to substrates such as histones (PubMed:28525742). During DNA repair, PARP1 recruits PARP9/BAL1-DTX3L complex to DNA damage sites via PARP9 binding to ribosylated PARP1 (PubMed:23230272). Subsequent PARP1-dependent PARP9/BAL1-DTX3L-mediated ubiquitination promotes the rapid and specific recruitment of 53BP1/TP53BP1, UIMC1/RAP80, and BRCA1 to DNA damage sites (PubMed:23230272, PubMed:28525742). In response to DNA damage, PARP9-DTX3L complex is required for efficient non-homologous end joining (NHEJ); the complex function is negatively modulated by PARP9 activity (PubMed:28525742). Dispensable for B-cell receptor (BCR) assembly through V(D)J recombination and class switch recombination (CSR) (By similarity). In macrophages, positively regulates pro-inflammatory cytokines production in response to IFNG stimulation by suppressing PARP14-mediated STAT1 ADP-ribosylation and thus promoting STAT1 phosphorylation (PubMed:27796300). Also suppresses PARP14-mediated STAT6 ADP-ribosylation (PubMed:27796300).[UniProtKB/Swiss-Prot Function]

Transcript Variant: This variant (1) encodes the longest isoform (a). Both variants 1 and 2 encode the same isoform (a).