

## Product datasheet for **SC314855**

### **PDPR (NM\_017990) Human Untagged Clone**

#### **Product data:**

<b>Product Type:</b>	Expression Plasmids
<b>Product Name:</b>	PDPR (NM_017990) Human Untagged Clone
<b>Tag:</b>	Tag Free
<b>Symbol:</b>	PDPR
<b>Synonyms:</b>	PDP3
<b>Mammalian Cell Selection:</b>	Neomycin
<b>Vector:</b>	pCMV6-Entry (PS100001)
<b>E. coli Selection:</b>	Kanamycin (25 ug/mL)
<b>Restriction Sites:</b>	Sgfl-MluI
<b>ACCN:</b>	NM_017990
<b>Insert Size:</b>	2640 bp
<b>OTI Disclaimer:</b>	<p>Due to the inherent nature of this plasmid, standard methods to replicate additional amounts of DNA in E. coli are highly likely to result in mutations and/or rearrangements. Therefore, OriGene does not guarantee the capability to replicate this plasmid DNA. Additional amounts of DNA can be purchased from OriGene with batch-specific, full-sequence verification at a reduced cost. Please contact our customer care team at <a href="mailto:custsupport@origene.com">custsupport@origene.com</a> or by calling 301.340.3188 option 3 for pricing and delivery.</p> <p>The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <a href="#">More info</a></p>
<b>OTI Annotation:</b>	This TrueClone is provided through our Custom Cloning Process that includes sub-cloning into OriGene's pCMV6 vector and full sequencing to provide a non-variant match to the expected reference without frameshifts, and is delivered as lyophilized plasmid DNA.
<b>Components:</b>	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).



[View online »](#)

**Reconstitution Method:**

1. Centrifuge at 5,000xg for 5min.
2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
3. Close the tube and incubate for 10 minutes at room temperature.
4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

**RefSeq:** [NM\\_017990.4](#)

**RefSeq Size:** 9255 bp

**RefSeq ORF:** 2640 bp

**Locus ID:** 55066

**UniProt ID:** [Q8NCN5](#)

**Cytogenetics:** 16q22.1

**Domains:** GCV\_T

**Protein Families:** Druggable Genome

**MW:** 99.4 kDa

**Gene Summary:** Pyruvate dehydrogenase complex (PDC) catalyzes the oxidative decarboxylation of pyruvate and links glycolysis to the tricarboxylic acid cycle and fatty acid synthesis. The dephosphorylation and reactivation of PDC is catalyzed by pyruvate dehydrogenase phosphatase (PDP). The dimeric PDP has a catalytic subunit and a regulatory subunit. This gene encodes the FAD-containing regulatory subunit of PDP. The encoded protein acts to decrease the sensitivity of the PDP catalytic subunit to magnesium ions. Alternative splicing results in multiple transcript variants encoding different isoforms. [provided by RefSeq, Jan 2017]