

Product datasheet for **SC315611**

PDE11A (NM_001077197) Human Untagged Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	PDE11A (NM_001077197) Human Untagged Clone
Tag:	Tag Free
Symbol:	PDE11A
Synonyms:	PPNAD2
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)




[View online »](#)

Fully Sequenced ORF: >SC315611 representing NM_001077197.
Blue=Insert sequence Red=Cloning site Green=Tag(s)

```
GCTCGTTTAGTGAACCGTCAGAATTTTGTAAACGACTCACTATAGGGCGGCCGGGAATTCGTGACTG
GATCCGGTACCGAGGAGATCTGCCGCCGCGATCGCC
ATGCTGAAGCAGGCAAGAAGACCTTTATTCAGAAATGTGCTCAGTGCCACACAGTGGAAAAAGTGAAA
ATCACAAGACTGGTCCAAATCTCTGGGGCCTCTTTGGTGAAAAACAGGAAAAGCACCAGGATTTTCTT
ATACAGAGGCAAACAAAAACAAAGGATCGACGATTCAATGATGAAATCGACAAGCTAACTGGATACAAG
ACAAAAATCATTATTGTGCATGCCTATCCGAAGCAGTGATGGTGAGATTATTGGTGTGCCCCAAGCGATA
AATAAGATTCTGAAGGAGCTCCATTTACTGAAGATGATGAAAAAGTTATGCAGATGTATCTTCCATTT
TGTGGAATCGCCATATCTAACGCTCAGCTCTTTGCTGCCTCAAGGAAAGAATATGAAAGAAGCAGAGCT
TTGCTAGAGGTGGTTAATGACCTCTTTGAAGAACAGACTGACCTGGAGAAAAATTGTCAAGAAAAAATG
CATCGGGCCAAACTCTGCTGAAATGTGAACGCTGTTCTGTTTTACTCCTAGAGGACATCGAATCACCA
GTGGTGAATTTACCAATCCTTTGAATTGATGTCCCAAAGTGCAGTGTGCTGATGCTGAGAACAGTTTC
AAAGAAAGCATGGAGAAATCATCACTCCGACTGGCTAATAAAACAGCATTGCTGAGCTGGTTGCT
TCAACAGGCCTTCCAGTGAACATCAGTGATGCCTACCAGGATCCGCGCTTTGATGCGAGGACAGACCAG
ATATCTGGTTTTACATAAGATCTGTTCTTTGTGTCCCTATTTGGAATAGCAACCACCAATAATTGGA
GTGGCTCAAGTGTAAACAGACTTGATGGGAAACCTTTTGATGATGCAGATCAACGACTTTTTGAGGCT
TTTGTCTCTTTTGTGGACTTGGCATCAACAACACAATTATGTATGATCAAGTGAAGAAGTCTGGGCC
AAGCAGTCTGTGGCTCTTGATGTGCTATCATACCATGCAACATGTTCAAAAGCTGAAGTTGACAAGTTT
AAGGCAGCCAACATCCCTCTGGTGCAGAACTTGCCATCGATGACATTCATTTTGATGACTTTTCTCTC
GACGTTGATGCCATGATCACAGCTGCTCTCCGGATGTTTATGGAGCTGGGGATGGTACAGAAATTTAAA
ATTGACTATGAGACACTGTGTAGGTGGCTTTTGACAGTGAGGAAAAACTATCGGATGGTTCTATACCAC
AACTGGAGACATGCCTTCAACGTGTGCAGCTGATGTTTCGCGATGTTAACCCTGCTGGGTTTCAAGAC
ATTCTGACCGAGGTGAAATTTTAGCGGTGATTGTGGGATGCCTGTGTCATGACCTCGACCACAGGGGA
ACCAACAATGCCTTCCAAGCTAAGAGTGGCTCTGCCCTGGCCAACTCTATGGAACCTCTGCTACCTTG
GAGCATCACCATTTCAACCACGCCGTGATGATCCTTCAAAGTGAGGGTCAACAATATCTTTGCTAACCTG
TCCTCAAAGGAATATAGTGACCTTATGCAGCTTTTGAAGCAGTCAATATTGGCAACAGACCTCACGCTG
TACTTTGAGAGGAGAAGTGAATTTTGAACCTGTGAGTAAAGGAGAATACGATTGGAACATCAAAAAC
CATCGTGATATATTTGATCAATGTTAATGACAGCCTGTGACCTTGGAGCCGTGACCAAACCGTGGGAG
ATCTCCAGACAGGTGGCAGAAGTTGTAACCAGTGAGTTCTTCGAACAAGGAGATCGGGAGAGATTAGAG
CTCAAACCTCACTCCTTACGAATTTTTGATCGGAACCGGAAGGATGAACTGCCTCGGTTGCAACTGGAG
TGGATTGATAGCATCTGCATGCCTTTGTATCAGGCACTGGTGAAGGTCAACGTGAAACTGAAGCCGATG
CTAGATTCACTAGCTACAAAACAGAAGTAAGTGGGAAGAGCTACACCAAAAACGACTGCTGGCCTCAACT
GCCTCATCCTCCCTGCCAGTGTATGGTAGCCAAGGAAGACAGGAACAA
ACGCGTACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAATGATATCCTGGAT
TACAAGGATGACGACGATAAGGTTTAAACGGCCGGC
```

Restriction Sites: Sgfl-Mlul

Plasmid Map:



ACCN: NM_001077197

Insert Size: 2052 bp

OTI Disclaimer: Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).

OTI Annotation: This TrueClone is provided through our Custom Cloning Process that includes sub-cloning into OriGene's pCMV6 vector and full sequencing to provide a non-variant match to the expected reference without frameshifts, and is delivered as lyophilized plasmid DNA.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method:

1. Centrifuge at 5,000xg for 5min.
2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
3. Close the tube and incubate for 10 minutes at room temperature.
4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

RefSeq: [NM_001077197.1](#)

RefSeq Size: 8316 bp

RefSeq ORF: 2052 bp

Locus ID: 50940

UniProt ID: [Q9HCR9](#)

Cytogenetics: 2q31.2

Protein Families:	Druggable Genome
Protein Pathways:	Progesterone-mediated oocyte maturation, Purine metabolism
MW:	78 kDa
Gene Summary:	<p>The 3',5'-cyclic nucleotides cAMP and cGMP function as second messengers in a wide variety of signal transduction pathways. 3',5'-cyclic nucleotide phosphodiesterases (PDEs) catalyze the hydrolysis of cAMP and cGMP to the corresponding 5'-monophosphates and provide a mechanism to downregulate cAMP and cGMP signaling. This gene encodes a member of the PDE protein superfamily. Mutations in this gene are a cause of Cushing disease and adrenocortical hyperplasia. Multiple transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Jul 2008]</p> <p>Transcript Variant: This variant (3) contains a distinct 5' UTR and 5' coding region, compared to variant 4. The resulting isoform (3) contains a shorter and distinct N-terminus, compared to isoform 4.</p>